

Final Examination.

Problem 4.

(30 marks)

Hand-written (complete) solution
by Prof. Lilian BESSON
26/11/2014.

Part A).

1) * For $p, k \in \mathbb{N}^*$, $p \geq 1$,so $x \mapsto \frac{1}{x^p}$ is decreasing because f is differentiable, and $f'(x) = \frac{-p}{x^{p+1}} < 0$.on $[1, +\infty)$ so on $[k, k+1]$ * Because f is decreasing on $[k, k+1]$.

$$\min_{[k, k+1]} f(x) = f(k+1) = \frac{1}{(k+1)^p}$$

$$\text{and } \max_{[k, k+1]} f(x) = f(k) = \frac{1}{k^p}$$

So the min-max inequality, applied to f .on $[k, k+1]$, gives

$$\frac{1}{(k+1)^p} \leq \int_k^{k+1} \frac{1}{t^p} dt \leq \frac{1}{k^p} \quad (1)$$

Problem 4

Part A

2) For $p \geq 1$, $\frac{1}{n^p}$ is well defined for all $n \geq 1$,
 so the finite sum $S_m(p) = \sum_{k=1}^m \frac{1}{k^p}$
 is well defined.

We know that

$$[1, m] = \bigcup_{k=1}^{m-1} [k, k+1]$$

So by summing the inequality (1)

for $k=1 \dots m-1$,

$$\sum_{k=1}^{m-1} \left(\frac{1}{k+1} \right)^p \leq \int_1^m f(x) dx \leq \sum_{k=1}^{m-1} \left(\frac{1}{k} \right)^p$$

$$\underline{S_m(p)} - 1 \leq \int_1^m f(x) dx \leq \overline{S_{m-1}(p)} -$$

3) For $A > 1$,

$$g(A) = \int_1^A f(x) dx = \int_1^A \frac{1}{x^p} dx$$

$$= \frac{1}{p-1} \left[\frac{1}{x^{p-1}} \right]_1^A = \frac{A^{1-p}}{p-1} - \frac{1}{p-1}$$

① If $p > 1$

so for $A \rightarrow +\infty$,

$$1-p < 1 \Rightarrow \frac{A^{1-p}}{p-1} \xrightarrow[A \rightarrow \infty]{} 0$$

$$\text{so } g(A) \rightarrow \frac{1}{1-p}$$

Problem 4

Part A

② If $p = 1$:
$$g(A) = \int_1^A \frac{1}{x} dx$$

$$= \ln(A) \xrightarrow[A \rightarrow +\infty]{} +\infty$$

Conclusion $\boxed{g(A) \text{ has a finite limit for } A \rightarrow +\infty \text{ if and only if } p > 2.}$

4). If $p \geq 2$ $g(n)$ has a finite limit
if when $n \rightarrow +\infty$,

so (2) gives:

$$\begin{aligned} g(n) &\leq S_{n-1}(p) = s_{n-1}(p) - 1 + 1 \\ &\leq g(n-1) + 1 \end{aligned}$$

But now $(g(n-1))_{n \in \mathbb{N}^*}$ is bounded because convergent
And $(S_m(p))_{m \in \mathbb{N}^*}$ is increasing and bounded

Conclusion $(S_m(p))_m$ converges.

If $p < 2$ ($\Rightarrow p = 1$) (2) gives

$$g(n) \leq S_{n-1}(p)$$

$$\downarrow n \rightarrow +\infty$$

$$\text{so } S_{n-1}(p) \xrightarrow[n \rightarrow +\infty]{} +\infty$$

Problem 4

Part A

4)

Conclusion. The sequence $S(p) = (S_n(p))_p$ is convergent if $p \geq 2$, divergent ($\rightarrow \infty$) if $p < 2$,
 so convergent if and only if $p \geq 2$.

Part B

$$1) \quad \varphi_o(t) = \begin{cases} -1 & \text{if } t = 0 \\ \frac{t^2/2\pi - t}{2\sin(t/2)} & \text{if } t \in (0, \pi] \end{cases}$$

Continuity: If $t \in (0, \pi]$, $\varphi_o(t)$ is continuous.
 Only $t=0$ is a potential problem.

$$\text{For } t \rightarrow 0^+, \quad \frac{\frac{t^2}{2\pi} - t}{2\sin(t/2)} = \frac{t/2}{\sin(t/2)} \times \left(\frac{t}{2\pi} - 1 \right)$$

and

$$\begin{cases} \frac{t}{2\pi} - 1 \xrightarrow[t \rightarrow 0^+]{} -1 \\ \frac{t/2}{\sin(t/2)} \xrightarrow[t \rightarrow 0^+]{} 1 \end{cases} \quad \text{so} \quad \boxed{\varphi_o(t) \xrightarrow[t \rightarrow 0^+]{} -1}$$

$$\varphi_o(0) \quad \boxed{-1}$$

So φ_o is indeed continuous on $[0, \pi]$.

Differentiability: For $t \in (0, \pi]$, φ_o is differentiable at t ,

$$\text{and } \varphi'_o(t) = \frac{\left(\frac{t}{\pi} - 1\right)}{2\sin(t/2)} - \frac{\left(\frac{t^2}{2\pi} - t\right) \cos(t/2)}{4\sin^2(t/2)}$$

$$\xrightarrow[t \rightarrow 0^+]{} \frac{1}{2\pi} \quad \text{if } t \rightarrow 0^+ \quad \text{so } \varphi'_o(0) \text{ exist.}$$