

# Recent progress on the Boltzmann equation without angular cutoff

Oral Defense

Julie Gauthier

University of Tsinghua, Department of Mathematical sciences

July, 15th, 2013

# Introduction : the Boltzmann equation without angular cutoff

$$\frac{\partial f}{\partial t} + \nabla_x f \cdot v = Q(f, f)$$

# Introduction : the Boltzmann equation without angular cutoff


$$\frac{\partial f}{\partial t} + \nabla_x f \cdot v = Q(f, f)$$

$$Q(g, f) = \int_{\mathbb{R}^N} \int_{S^{N-1}} B(g'_* f' - g_* f) d\sigma dv_*$$

with

$$v' = \frac{v + v_*}{2} + \frac{|v - v_*|}{2} \sigma$$

$$v'_* = \frac{v + v_*}{2} - \frac{|v - v_*|}{2} \sigma$$



$$\begin{aligned} B(v - v_*, \sigma) &= B(|v - v_*|, k \cdot \sigma) \\ &= B(|v - v_*|, \cos \theta) \end{aligned}$$

# Introduction : the Boltzmann equation without angular cutoff


$$\frac{\partial f}{\partial t} + \nabla_x f \cdot v = Q(f, f)$$

$$Q(g, f) = \int_{\mathbb{R}^N} \int_{S^{N-1}} B(g'_* f' - g_* f) d\sigma dv_*$$

with

$$v' = \frac{v + v_*}{2} + \frac{|v - v_*|}{2} \sigma$$

$$v'_* = \frac{v + v_*}{2} - \frac{|v - v_*|}{2} \sigma$$



Non-cutoff assumption

$$\int_{S^{N-1}} B(z, \sigma) d\sigma = +\infty$$

$$\begin{aligned} B(v - v_*, \sigma) &= B(|v - v_*|, k \cdot \sigma) \\ &= B(|v - v_*|, \cos \theta) \end{aligned}$$

# Introduction : the Boltzmann equation without angular cutoff

$$\frac{\partial f}{\partial t} + \nabla_x f \cdot v = Q(f, f)$$


$$Q(g, f) = \int_{\mathbb{R}^N} \int_{S^{N-1}} B(g'_* f' - g_* f) d\sigma dv_*$$

with

$$v' = \frac{v + v_*}{2} + \frac{|v - v_*|}{2} \sigma$$

$$v'_* = \frac{v + v_*}{2} - \frac{|v - v_*|}{2} \sigma$$

$$\begin{aligned} B(v - v_*, \sigma) &= B(|v - v_*|, k \cdot \sigma) \\ &= B(|v - v_*|, \cos \theta) \end{aligned}$$



Non-cutoff assumption

$$\int_{S^{N-1}} B(z, \sigma) d\sigma = +\infty$$

$$B(z, \sigma) \geq \Phi(|z|) b(\cos \theta)$$

with  $\Phi(|z|) = |z|^\gamma$  and

$$\sin^{N-2} \theta b(\cos \theta) \underset{\theta \rightarrow 0^+}{\sim} \frac{K}{\theta^{1+\nu}}$$

# *Entropy Dissipation and Long-Range Interactions*

Main result

$$\|\sqrt{f}\|_{H^{\nu/2}(|v| < R)}^2 \leq C_{g,R} \left( D(g, f) + \|g\|_{L_2^1} \|f\|_{L_2^1} \right)$$

# *Entropy Dissipation and Long-Range Interactions*

Main result

$$\|\sqrt{f}\|_{H^{\nu/2}(|v| < R)}^2 \leq C_{g,R} \left( D(g, f) + \|g\|_{L_2^1} \|f\|_{L_2^1} \right)$$

$$D(g, f) = - \int_{\mathbb{R}^N} Q(g, f) \log(f) dv$$

# Entropy Dissipation and Long-Range Interactions

## Main result

$$\|\sqrt{f}\|_{H^{\nu/2}(|v| < R)}^2 \leq C_{g,R} \left( D(g, f) + \|g\|_{L_2^1} \|f\|_{L_2^1} \right)$$

$$D(g, f) = - \int_{\mathbb{R}^N} Q(g, f) \log(f) dv$$

## Assumptions on $B$

- non-cutoff ( $\Phi$  continuous, positive except in 0,  $0 \leq \nu < 2$ )
- $\Lambda(|z|) + |z|\Lambda'(|z|) \leq C_0(1 + |z|)^2$   
where

$$\Lambda(|z|) = \int_{S^{N-1}} B(z, \sigma) (1 - k \cdot \sigma) d\sigma$$

and

$$\Lambda'(|z|) = \int_{S^{N-1}} B'(z, \sigma) (1 - k \cdot \sigma) d\sigma$$

with  $B'(z, \sigma) =$

$$\sup_{1 < \lambda \leq \sqrt{2}} \frac{|B(\lambda z, \sigma) - B(z, \sigma)|}{(\lambda - 1)|z|}.$$

# Entropy Dissipation and Long-Range Interactions

## Main result

$$\|\sqrt{f}\|_{H^{\nu/2}(|v| < R)}^2 \leq C_{g,R} \left( D(g, f) + \|g\|_{L_2^1} \|f\|_{L_2^1} \right)$$

$$D(g, f) = - \int_{\mathbb{R}^N} Q(g, f) \log(f) dv$$

## Assumptions on $B$

## Proof step by step

- non-cutoff ( $\Phi$  continuous, positive except in 0,  $0 \leq \nu < 2$ )
- $\Lambda(|z|) + |z|\Lambda'(|z|) \leq C_0(1 + |z|)^2$   
where

$$\Lambda(|z|) = \int_{S^{N-1}} B(z, \sigma) (1 - k \cdot \sigma) d\sigma$$

and

$$\Lambda'(|z|) = \int_{S^{N-1}} B'(z, \sigma) (1 - k \cdot \sigma) d\sigma$$

with  $B'(z, \sigma) =$

$$\sup_{1 < \lambda \leq \sqrt{2}} \frac{|B(\lambda z, \sigma) - B(z, \sigma)|}{(\lambda - 1)|z|}.$$

# Entropy Dissipation and Long-Range Interactions

## Main result

$$\|\sqrt{f}\|_{H^{\nu/2}(|v| < R)}^2 \leq C_{g,R} \left( D(g, f) + \|g\|_{L_2^1} \|f\|_{L_2^1} \right)$$

$$D(g, f) = - \int_{\mathbb{R}^N} Q(g, f) \log(f) dv$$

## Assumptions on $B$

- non-cutoff ( $\Phi$  continuous, positive except in 0,  $0 \leq \nu < 2$ )
- $\Lambda(|z|) + |z|\Lambda'(|z|) \leq C_0(1 + |z|)^2$  where

$$\Lambda(|z|) = \int_{S^{N-1}} B(z, \sigma) (1 - k \cdot \sigma) d\sigma$$

and

$$\Lambda'(|z|) = \int_{S^{N-1}} B'(z, \sigma) (1 - k \cdot \sigma) d\sigma$$

with  $B'(z, \sigma) =$

$$\sup_{1 < \lambda \leq \sqrt{2}} \frac{|B(\lambda z, \sigma) - B(z, \sigma)|}{(\lambda - 1)|z|}.$$

## Proof step by step

- cut  $D(g, f)$  into two parts by habile use of two changes of variables

# Entropy Dissipation and Long-Range Interactions

## Main result

$$\|\sqrt{f}\|_{H^{\nu/2}(|v| < R)}^2 \leq C_{g,R} \left( D(g, f) + \|g\|_{L_2^1} \|f\|_{L_2^1} \right)$$

$$D(g, f) = - \int_{\mathbb{R}^N} Q(g, f) \log(f) dv$$

## Assumptions on $B$

- non-cutoff ( $\Phi$  continuous, positive except in 0,  $0 \leq \nu < 2$ )

- $\Lambda(|z|) + |z|\Lambda'(|z|) \leq C_0(1 + |z|)^2$   
where

$$\Lambda(|z|) = \int_{S^{N-1}} B(z, \sigma)(1 - k \cdot \sigma) d\sigma$$

and

$$\Lambda'(|z|) = \int_{S^{N-1}} B'(z, \sigma)(1 - k \cdot \sigma) d\sigma$$

with  $B'(z, \sigma) =$

$$\sup_{1 < \lambda \leq \sqrt{2}} \frac{|B(\lambda z, \sigma) - B(z, \sigma)|}{(\lambda - 1)|z|}.$$

## Proof step by step

- cut  $D(g, f)$  into two parts by habile use of two changes of variables
- bound the first part  
 $\implies$  cancellation lemma

# Entropy Dissipation and Long-Range Interactions

## Main result

$$\|\sqrt{f}\|_{H^{\nu/2}(|v| < R)}^2 \leq C_{g,R} \left( D(g, f) + \|g\|_{L_2^1} \|f\|_{L_2^1} \right)$$

$$D(g, f) = - \int_{\mathbb{R}^N} Q(g, f) \log(f) dv$$

## Assumptions on $B$

- non-cutoff ( $\Phi$  continuous, positive except in 0,  $0 \leq \nu < 2$ )

- $\Lambda(|z|) + |z|\Lambda'(|z|) \leq C_0(1 + |z|)^2$  where

$$\Lambda(|z|) = \int_{S^{N-1}} B(z, \sigma)(1 - k \cdot \sigma) d\sigma \text{ and}$$

$$\Lambda'(|z|) = \int_{S^{N-1}} B'(z, \sigma)(1 - k \cdot \sigma) d\sigma \text{ with } B'(z, \sigma) =$$

$$\sup_{1 < \lambda \leq \sqrt{2}} \frac{|B(\lambda z, \sigma) - B(z, \sigma)|}{(\lambda - 1)|z|}.$$

## Proof step by step

- cut  $D(g, f)$  into two parts by habile use of two changes of variables
- bound the first part  
 $\implies$  cancellation lemma
- first assume  $\Phi = 1$  and use Fourier analysis to bound the second part

# Entropy Dissipation and Long-Range Interactions

## Main result

$$\|\sqrt{f}\|_{H^{\nu/2}(|v| < R)}^2 \leq C_{g,R} \left( D(g, f) + \|g\|_{L_2^1} \|f\|_{L_2^1} \right)$$

$$D(g, f) = - \int_{\mathbb{R}^N} Q(g, f) \log(f) dv$$

## Assumptions on $B$

- non-cutoff ( $\Phi$  continuous, positive except in 0,  $0 \leq \nu < 2$ )
- $\Lambda(|z|) + |z|\Lambda'(|z|) \leq C_0(1 + |z|)^2$  where

$$\Lambda(|z|) = \int_{S^{N-1}} B(z, \sigma)(1 - k \cdot \sigma) d\sigma \quad \text{and}$$

$$\Lambda'(|z|) = \int_{S^{N-1}} B'(z, \sigma)(1 - k \cdot \sigma) d\sigma \quad \text{with } B'(z, \sigma) = \sup_{1 < \lambda \leq \sqrt{2}} \frac{|B(\lambda z, \sigma) - B(z, \sigma)|}{(\lambda - 1)|z|}.$$

## Proof step by step

- cut  $D(g, f)$  into two parts by habile use of two changes of variables
- bound the first part  
 $\implies$  cancellation lemma
- first assume  $\Phi = 1$  and use Fourier analysis to bound the second part
- generalize

## Cancellation lemma

$$D(g, f) = \int Bg_*(f - f') d\nu d\nu_* d\sigma + \int Bg_* \left( f \log \frac{f}{f'} - f + f' \right) d\nu d\nu_* d\sigma$$

## Cancellation lemma

$$D(g, f) = \int Bg_*(f - f') d\nu d\nu_* d\sigma + \int Bg_* \left( f \log \frac{f}{f'} - f + f' \right) d\nu d\nu_* d\sigma$$

Theorem (Cancellation lemma)

$$S(z) = |S^{N-2}| \int_0^{\frac{\pi}{2}} \sin^{N-2} \theta \left( \frac{B\left(\frac{|z|}{\cos \frac{\theta}{2}}, \cos \theta\right)}{\cos^{N-2} \theta} - B(|z|, \cos \theta) \right) d\theta$$

We have:

1.  $|S(z)| \leq C_N (\Lambda(|z|) + |z|\Lambda'(|z|))$
2.  $\int B(f' - f) d\nu d\sigma = (f \star S)(\nu_*)$

## Cancellation lemma

$$D(g, f) = \int Bg_*(f - f')dv dv_* d\sigma + \int Bg_* \left( f \log \frac{f}{f'} - f + f' \right) dv dv_* d\sigma$$

Theorem (Cancellation lemma)

$$S(z) = |S^{N-2}| \int_0^{\frac{\pi}{2}} \sin^{N-2} \theta \left( \frac{B\left(\frac{|z|}{\cos \frac{\theta}{2}}, \cos \theta\right)}{\cos^N \frac{\theta}{2}} - B(|z|, \cos \theta) \right) d\theta$$

We have:

1.  $|S(z)| \leq C_N (\Lambda(|z|) + |z|\Lambda'(|z|))$
2.  $\int B(f' - f)dv d\sigma = (f \star S)(v_*)$

## Corollary

There exists  $C$  depending only on  $N$  and  $B$   
such that

$$|\int Bg_*(f' - f)dv dv_* d\sigma| \leq C \|g\|_{L^1_2} \|f\|_{L^1_2}$$

## Cancellation lemma

$$D(g, f) = \int Bg_*(f - f')dv dv_* d\sigma + \int Bg_* \left( f \log \frac{f}{f'} - f + f' \right) dv dv_* d\sigma$$

Theorem (Cancellation lemma)

$$S(z) = |S^{N-2}| \int_0^{\frac{\pi}{2}} \sin^{N-2} \theta \left( \frac{B\left(\frac{|z|}{\cos \frac{\theta}{2}}, \cos \theta\right)}{\cos^{N-2} \theta} - B(|z|, \cos \theta) \right) d\theta$$

We have:

1.  $|S(z)| \leq C_N (\Lambda(|z|) + |z|\Lambda'(|z|))$
2.  $\int B(f' - f)dv d\sigma = (f \star S)(v_*)$

## Corollary

There exists  $C$  depending only on  $N$  and  $B$  such that

$$|\int Bg_*(f' - f)dv dv_* d\sigma| \leq C \|g\|_{L^1_2} \|f\|_{L^1_2}$$

## Proof.

Use of the change of variables  $v \rightarrow v'$  forth and backward. □

Use of *Fourier Analysis* for the second term, easy case :

$$\Phi = 1$$

$$\int B g_* \left( f \log \frac{f}{F'} - f + f' \right) d\nu d\nu_* d\sigma \geq \int \Phi b g_* (F' - F)^2 d\nu d\nu_* d\sigma$$

Use of *Fourier Analysis* for the second term, easy case :

$$\Phi = 1$$

$$\int B g_* \left( f \log \frac{f}{F'} - f + f' \right) dv dv_* d\sigma \geq \int \Phi b g_*(F' - F)^2 dv dv_* d\sigma$$

Fourier transform  $\implies$  with  $\xi^- = \frac{\xi - |\xi|\sigma}{2}$ ,

$$\geq \frac{1}{(2\pi)^N} \int \left| \hat{F}(\xi) \right|^2 \left( \int b \left( \frac{\xi}{|\xi|} \cdot \sigma \right) (\hat{g}(0) - |\hat{g}(\xi^-)|) d\sigma \right) d\xi$$


Use of Fourier Analysis for the second term, easy case :

$$\Phi = 1$$

$$\int B g_* \left( f \log \frac{f}{F'} - f + f' \right) dv dv_* d\sigma \geq \int \Phi b g_*(F' - F)^2 dv dv_* d\sigma$$

Fourier transform  $\implies$  with  $\xi^- = \frac{\xi - |\xi|\sigma}{2}$ ,

$$\geq \frac{1}{(2\pi)^N} \int \left| \hat{F}(\xi) \right|^2 \left( \int b \left( \frac{\xi}{|\xi|} \cdot \sigma \right) (\hat{g}(0) - |\hat{g}(\xi^-)|) d\sigma \right) d\xi$$



$$A = \{ |v| \leq r, \exists p \in \mathbb{Z}, |v \cdot \xi^- + \alpha - 2p\pi| < 2\varepsilon \}$$

$$\hat{g}(0) - |\hat{g}(\xi^-)| = 2 \int g(v) \sin^2 \left( \frac{v \cdot \xi^- + \alpha}{2} \right) dv$$

$$\geq 2 \sin^2 \varepsilon \left( \|g\|_{L_1^1(\mathbb{R}^N)} - \frac{\|g\|_{L_1^1(\mathbb{R}^N)}}{r} - \int_A g(v) dv \right)$$


Use of Fourier Analysis for the second term, easy case :

$$\Phi = 1$$

$$\int B g_* \left( f \log \frac{f}{f'} - f + f' \right) d\nu d\nu_* d\sigma \geq \int \Phi b g_*(F' - F)^2 d\nu d\nu_* d\sigma$$

Fourier transform  $\implies$  with  $\xi^- = \frac{\xi - |\xi|\sigma}{2}$ ,

$$\geq \frac{1}{(2\pi)^N} \int \left| \hat{F}(\xi) \right|^2 \left( \int b \left( \frac{\xi}{|\xi|} \cdot \sigma \right) (\hat{g}(0) - |\hat{g}(\xi^-)|) d\sigma \right) d\xi$$



$$A = \{ |v| \leq r, \exists p \in \mathbb{Z}, |v \cdot \xi^- + \alpha - 2p\pi| < 2\varepsilon \}$$

$$\hat{g}(0) - |\hat{g}(\xi^-)| = 2 \int g(v) \sin^2 \left( \frac{v \cdot \xi^- + \alpha}{2} \right) dv$$

$$\geq 2 \sin^2 \varepsilon \left( \|g\|_{L_1^1(\mathbb{R}^N)} - \frac{\|g\|_{L_1^1(\mathbb{R}^N)}}{r} - \int_A g(v) dv \right)$$

$$\geq 2 \sin^2 \varepsilon \left( \|g\|_{L_1^1(\mathbb{R}^N)} - \frac{\|g\|_{L_1^1(\mathbb{R}^N)}}{r} - \sup_{|A| \leq C_{\varepsilon, r, \xi^-}} \int_A g(v) dv \right)$$

# Truncature : generalization

## Problems

1.  $\Phi(|z|) \rightarrow \infty$
2.  $\Phi(0) = 0$

# Truncature : generalization

## Problems

1.  $\Phi(|z|) \rightarrow \infty$
2.  $\Phi(0) = 0$

## Idea

$f \rightarrow f_A, g \rightarrow g_B$

# Truncature : generalization

## Problems

1.  $\Phi(|z|) \rightarrow \infty$
2.  $\Phi(0) = 0$

## Idea

$f \rightarrow f_A, g \rightarrow g_B$

## First case

$A = B = B_R$

# Truncature : generalization

## Second case

Problems


1.  $\Phi(|z|) \rightarrow \infty$
2.  $\Phi(0) = 0$

Idea

$$f \rightarrow f_A, g \rightarrow g_B$$

First case

$$A = B = B_R$$



# *On the Boltzmann Equation for Long-Range Interactions*

Existence of weak solutions, appearance of strong compactness.

# *On the Boltzmann Equation for Long-Range Interactions*

Existence of weak solutions, appearance of strong compactness.

## **Definition (weak solution)**

*f: renormalized solution to the Boltzmann equation with defect measure*

$$f \in C(\mathbb{R}^+; \mathcal{D}'(\mathbb{R}_x^N \times \mathbb{R}_v^N)) \cap L^\infty(\mathbb{R}^+; L^1((1 + |v|^2 + |x|^2) dx dv))$$

$\forall \beta \in C^2(\mathbb{R}^+, \mathbb{R}^+)$  non-linear such that :

$$\begin{aligned} \beta(0) = 0 && 0 < \beta'(f) \leq \frac{C}{1+f} && \beta''(f) < 0 \end{aligned}$$

$$\implies \begin{cases} \frac{\partial \beta(f)}{\partial t} + v \cdot \nabla_x \beta(f) \geq \beta'(f) Q(f, f) \\ \forall t \geq 0, \int_{\mathbb{R}^N \times \mathbb{R}^N} f(t, x, v) dx dv = \int_{\mathbb{R}^N \times \mathbb{R}^N} f(0, x, v) dx dv \end{cases}$$

## Assumptions on $B$

AT MOST BORDERLINE

SINGULARITY.

$$B(z, \sigma) = \frac{\beta_0(k \cdot \sigma)}{|z|^N} + B_1(z, \sigma) \text{ with}$$

- $\int_{S^{N-1}} \beta_0(k \cdot \sigma)(1 - k \cdot \sigma) d\sigma < +\infty$
- $\Lambda_1(|z|), |z|\Lambda'_1(|z|) \in L^1_{\text{loc}}(\mathbb{R}^N)$

## Assumptions on $B$

AT MOST BORDERLINE

SINGULARITY.

$$B(z, \sigma) = \frac{\beta_0(k.\sigma)}{|z|^N} + B_1(z, \sigma) \text{ with}$$

- $\int_{S^{N-1}} \beta_0(k.\sigma)(1 - k.\sigma) d\sigma < +\infty$
- $\Lambda_1(|z|), |z|\Lambda'_1(|z|) \in L^1_{\text{loc}}(\mathbb{R}^N)$

BEHAVIOUR AT INFINITY.

$$M^\alpha(|z|) := \int_{S^{N-1}} B(z, \sigma) (1 - k.\sigma)^{\frac{\alpha}{2}} d\sigma$$

- $\exists \alpha \in [0 ; 2], M^\alpha(|z|) = \underset{|z| \rightarrow \infty}{o} (|z|^{2-\alpha})$
- $|z|M'(|z|) = \underset{|z| \rightarrow \infty}{o} (|z|^2)$

## Assumptions on $B$

AT MOST BORDERLINE

SINGULARITY.

$$B(z, \sigma) = \frac{\beta_0(k.\sigma)}{|z|^N} + B_1(z, \sigma) \text{ with}$$

- $\int_{S^{N-1}} \beta_0(k.\sigma)(1 - k.\sigma) d\sigma < +\infty$
- $\Lambda_1(|z|), |z|\Lambda'_1(|z|) \in L^1_{\text{loc}}(\mathbb{R}^N)$

BEHAVIOUR AT INFINITY.

$$M^\alpha(|z|) := \int_{S^{N-1}} B(z, \sigma)(1 - k.\sigma)^{\frac{\alpha}{2}} d\sigma$$

- $\exists \alpha \in [0 ; 2], M^\alpha(|z|) = \underset{|z| \rightarrow \infty}{o} (|z|^{2-\alpha})$
- $|z|M'(|z|) = \underset{|z| \rightarrow \infty}{o} (|z|^2)$

ANGULAR SINGULARITY

CONDITION.  $\int_{S^{N-1}} b(k.\sigma) d\sigma = +\infty$

## Assumptions on $B$

AT MOST BORDERLINE  
SINGULARITY.

$$B(z, \sigma) = \frac{\beta_0(k.\sigma)}{|z|^N} + B_1(z, \sigma) \text{ with}$$

- $\int_{S^{N-1}} \beta_0(k.\sigma)(1 - k.\sigma) d\sigma < +\infty$
- $\Lambda_1(|z|), |z|\Lambda'_1(|z|) \in L^1_{\text{loc}}(\mathbb{R}^N)$

BEHAVIOUR AT INFINITY.

$$M^\alpha(|z|) := \int_{S^{N-1}} B(z, \sigma)(1 - k.\sigma)^{\frac{\alpha}{2}} d\sigma$$

- $\exists \alpha \in [0 ; 2], M^\alpha(|z|) = \underset{|z| \rightarrow \infty}{o} (|z|^{2-\alpha})$
- $|z|M'(|z|) = \underset{|z| \rightarrow \infty}{o} (|z|^2)$

ANGULAR SINGULARITY

CONDITION.  $\int_{S^{N-1}} b(k.\sigma) d\sigma = +\infty$

$B^n$  APPROXIMATES  $B$ .

- $S^n \rightharpoonup S$  locally in the weak-measure sense

## Assumptions on $B$

AT MOST BORDERLINE  
SINGULARITY.

$$B(z, \sigma) = \frac{\beta_0(k.\sigma)}{|z|^N} + B_1(z, \sigma) \text{ with}$$

- $\int_{S^{N-1}} \beta_0(k.\sigma)(1 - k.\sigma) d\sigma < +\infty$
- $\Lambda_1(|z|), |z|\Lambda'_1(|z|) \in L^1_{\text{loc}}(\mathbb{R}^N)$

BEHAVIOUR AT INFINITY.

$$M^\alpha(|z|) := \int_{S^{N-1}} B(z, \sigma)(1 - k.\sigma)^{\frac{\alpha}{2}} d\sigma$$

- $\exists \alpha \in [0 ; 2], M^\alpha(|z|) = \underset{|z| \rightarrow \infty}{o} (|z|^{2-\alpha})$
- $|z|M'(|z|) = \underset{|z| \rightarrow \infty}{o} (|z|^2)$

ANGULAR SINGULARITY

CONDITION.  $\int_{S^{N-1}} b(k.\sigma) d\sigma = +\infty$

$B^n$  APPROXIMATES  $B$ .

- $S^n \rightharpoonup S$  locally in the weak-measure sense
- For  $\mathcal{T} : \phi \mapsto \int_{S^{N-1}} B(\phi' - \phi) d\sigma$ ,  
 $\mathcal{T}^n \rightharpoonup \mathcal{T}$

## Assumptions on $B$

AT MOST BORDERLINE  
SINGULARITY.

$$B(z, \sigma) = \frac{\beta_0(k.\sigma)}{|z|^N} + B_1(z, \sigma) \text{ with}$$

- $\int_{S^{N-1}} \beta_0(k.\sigma)(1 - k.\sigma) d\sigma < +\infty$
- $\Lambda_1(|z|), |z|\Lambda'_1(|z|) \in L^1_{loc}(\mathbb{R}^N)$

BEHAVIOUR AT INFINITY.

$$M^\alpha(|z|) := \int_{S^{N-1}} B(z, \sigma)(1 - k.\sigma)^{\frac{\alpha}{2}} d\sigma$$

- $\exists \alpha \in [0 ; 2], M^\alpha(|z|) = \underset{|z| \rightarrow \infty}{o} (|z|^{2-\alpha})$
- $|z|M'(|z|) = \underset{|z| \rightarrow \infty}{o} (|z|^2)$

ANGULAR SINGULARITY

CONDITION.  $\int_{S^{N-1}} b(k.\sigma) d\sigma = +\infty$

$B^n$  APPROXIMATES  $B$ .

- $S^n \rightharpoonup S$  locally in the weak-measure sense
- For  $\mathcal{T} : \phi \mapsto \int_{S^{N-1}} B(\phi' - \phi) d\sigma$ ,  
 $\mathcal{T}^n \rightharpoonup \mathcal{T}$
- $B^n \rightarrow B$  a.e.

## Assumptions on $B$

AT MOST BORDERLINE  
SINGULARITY.

$$B(z, \sigma) = \frac{\beta_0(k.\sigma)}{|z|^N} + B_1(z, \sigma) \text{ with}$$

- $\int_{S^{N-1}} \beta_0(k.\sigma)(1 - k.\sigma) d\sigma < +\infty$
- $\Lambda_1(|z|), |z|\Lambda'_1(|z|) \in L^1_{\text{loc}}(\mathbb{R}^N)$

BEHAVIOUR AT INFINITY.

$$M^\alpha(|z|) := \int_{S^{N-1}} B(z, \sigma)(1 - k.\sigma)^{\frac{\alpha}{2}} d\sigma$$

- $\exists \alpha \in [0 ; 2], M^\alpha(|z|) = \underset{|z| \rightarrow \infty}{o} (|z|^{2-\alpha})$
- $|z|M'(|z|) = \underset{|z| \rightarrow \infty}{o} (|z|^2)$

ANGULAR SINGULARITY

CONDITION.  $\int_{S^{N-1}} b(k.\sigma) d\sigma = +\infty$

$B^n$  APPROXIMATES  $B$ .

- $S^n \rightharpoonup S$  locally in the weak-measure sense
- For  $\mathcal{T} : \phi \mapsto \int_{S^{N-1}} B(\phi' - \phi) d\sigma$ ,  
 $\mathcal{T}^n \rightharpoonup \mathcal{T}$
- $B^n \rightarrow B$  a.e.
- BEHAVIOUR AT INFINITY uniformly in  $n$ .

## Assumptions on $B$

AT MOST BORDERLINE  
SINGULARITY.

$$B(z, \sigma) = \frac{\beta_0(k.\sigma)}{|z|^N} + B_1(z, \sigma) \text{ with}$$

- $\int_{S^{N-1}} \beta_0(k.\sigma)(1 - k.\sigma) d\sigma < +\infty$
- $\Lambda_1(|z|), |z|\Lambda'_1(|z|) \in L^1_{\text{loc}}(\mathbb{R}^N)$

BEHAVIOUR AT INFINITY.

$$M^\alpha(|z|) := \int_{S^{N-1}} B(z, \sigma)(1 - k.\sigma)^{\frac{\alpha}{2}} d\sigma$$

- $\exists \alpha \in [0 ; 2], M^\alpha(|z|) = \underset{|z| \rightarrow \infty}{o} (|z|^{2-\alpha})$
- $|z|M'(|z|) = \underset{|z| \rightarrow \infty}{o} (|z|^2)$

ANGULAR SINGULARITY  
CONDITION.  $\int_{S^{N-1}} b(k.\sigma) d\sigma = +\infty$

$B^n$  APPROXIMATES  $B$ .

- $S^n \rightharpoonup S$  locally in the weak-measure sense
- For  $\mathcal{T} : \phi \mapsto \int_{S^{N-1}} B(\phi' - \phi) d\sigma$ ,  
 $\mathcal{T}^n \rightharpoonup \mathcal{T}$
- $B^n \rightarrow B$  a.e.
- BEHAVIOUR AT INFINITY uniformly in  $n$ .

OVERALL ANGULAR SINGULARITY  
CONDITION.

## Assumptions on $B$

AT MOST BORDERLINE SINGULARITY.

$$B(z, \sigma) = \frac{\beta_0(k.\sigma)}{|z|^N} + B_1(z, \sigma) \text{ with}$$

- $\int_{S^{N-1}} \beta_0(k.\sigma)(1 - k.\sigma) d\sigma < +\infty$
- $\Lambda_1(|z|), |z|\Lambda'_1(|z|) \in L^1_{loc}(\mathbb{R}^N)$

BEHAVIOUR AT INFINITY.

$$M^\alpha(|z|) := \int_{S^{N-1}} B(z, \sigma)(1 - k.\sigma)^{\frac{\alpha}{2}} d\sigma$$

- $\exists \alpha \in [0 ; 2], M^\alpha(|z|) = \underset{|z| \rightarrow \infty}{o} (|z|^{2-\alpha})$
- $|z|M'(|z|) = \underset{|z| \rightarrow \infty}{o} (|z|^2)$

ANGULAR SINGULARITY CONDITION.  $\int_{S^{N-1}} b(k.\sigma) d\sigma = +\infty$

$B^n$  APPROXIMATES  $B$ .

- $S^n \rightharpoonup S$  locally in the weak-measure sense
- For  $\mathcal{T} : \phi \mapsto \int_{S^{N-1}} B(\phi' - \phi) d\sigma$ ,  
 $\mathcal{T}^n \rightharpoonup \mathcal{T}$
- $B^n \rightarrow B$  a.e.
- BEHAVIOUR AT INFINITY uniformly in  $n$ .

OVERALL ANGULAR SINGULARITY CONDITION.

- $\forall n$ , ANGULAR SINGULARITY CONDITION

## Assumptions on $B$

AT MOST BORDERLINE SINGULARITY.

$$B(z, \sigma) = \frac{\beta_0(k.\sigma)}{|z|^N} + B_1(z, \sigma) \text{ with}$$

- $\int_{S^{N-1}} \beta_0(k.\sigma)(1 - k.\sigma) d\sigma < +\infty$
- $\Lambda_1(|z|), |z|\Lambda'_1(|z|) \in L^1_{\text{loc}}(\mathbb{R}^N)$

BEHAVIOUR AT INFINITY.

$$M^\alpha(|z|) := \int_{S^{N-1}} B(z, \sigma)(1 - k.\sigma)^{\frac{\alpha}{2}} d\sigma$$

- $\exists \alpha \in [0 ; 2], M^\alpha(|z|) = \underset{|z| \rightarrow \infty}{o} (|z|^{2-\alpha})$
- $|z|M'(|z|) = \underset{|z| \rightarrow \infty}{o} (|z|^2)$

ANGULAR SINGULARITY CONDITION.  $\int_{S^{N-1}} b(k.\sigma) d\sigma = +\infty$

$B^n$  APPROXIMATES  $B$ .

- $S^n \rightharpoonup S$  locally in the weak-measure sense
- For  $\mathcal{T} : \phi \mapsto \int_{S^{N-1}} B(\phi' - \phi) d\sigma$ ,  
 $\mathcal{T}^n \rightharpoonup \mathcal{T}$
- $B^n \rightarrow B$  a.e.
- BEHAVIOUR AT INFINITY uniformly in  $n$ .

OVERALL ANGULAR SINGULARITY CONDITION.

- $\forall n$ , ANGULAR SINGULARITY CONDITION
- $\int_{S^{N-1}} \lim_{n \rightarrow \infty} b^n(k.\sigma) d\sigma = \infty$

## Idea of the proof

$\beta'(f)Q(f, f)$  formally cut into three parts :

- something  $\times \int B(f'_* - f_*)dv_*d\sigma \implies$  use of the *cancellation lemma*
- $Q(f, \beta(f)) \implies$  apparition of  $\mathcal{T}$
- $-\int Bf'_*\Gamma(f, f')dv_*d\sigma$  which convergence is a consequence of the convergence of the first two terms

$$\Gamma(f, f') := \beta(f') - \beta(f) - \beta'(f)(f' - f) \leq 0$$

## Computation of $S$

$$S(|z|) = \lambda\delta_0 + S_1(|z|)$$

## Computation of $S$

$$S(|z|) = \lambda\delta_0 + S_1(|z|)$$

**Proof.**

$B_\varepsilon := \frac{\beta_0}{|z|^N} \mathbf{1}_{|z| \geq \varepsilon}$  gives  $S_\varepsilon$  which goes to  $S_0$  when  $\varepsilon \rightarrow 0$

$$\begin{aligned} S_\varepsilon(|z|) &\propto \frac{1}{|z|^N} \int_0^{\frac{\pi}{2}} \sin^{N-2} \theta \left( \mathbf{1}_{|z| \geq \varepsilon \cos \frac{\theta}{2}} - \mathbf{1}_{|z| \geq \varepsilon} \right) \beta_0(\cos \theta) d\theta \\ &\propto \frac{1}{\varepsilon^N} \frac{1}{(|z|/\varepsilon)^N} \int_{2 \cos^{-1} \frac{|z|}{\varepsilon}}^{\frac{\pi}{2}} \sin^{N-2} \theta \beta_0(\cos \theta) d\theta \mathbf{1}_{\frac{|z|}{\varepsilon} \leq 1} \\ &\propto \frac{1}{\varepsilon^N} J\left(\frac{|z|}{\varepsilon}\right) \end{aligned}$$

## Computation of $S$

$$S(|z|) = \lambda\delta_0 + S_1(|z|)$$

**Proof.**

$B_\varepsilon := \frac{\beta_0}{|z|^N} \mathbf{1}_{|z| \geq \varepsilon}$  gives  $S_\varepsilon$  which goes to  $S_0$  when  $\varepsilon \rightarrow 0$

$$\begin{aligned} S_\varepsilon(|z|) &\propto \frac{1}{|z|^N} \int_0^{\frac{\pi}{2}} \sin^{N-2} \theta \left( \mathbf{1}_{|z| \geq \varepsilon \cos \frac{\theta}{2}} - \mathbf{1}_{|z| \geq \varepsilon} \right) \beta_0(\cos \theta) d\theta \\ &\propto \frac{1}{\varepsilon^N} \frac{1}{(|z|/\varepsilon)^N} \int_{2 \cos^{-1} \frac{|z|}{\varepsilon}}^{\frac{\pi}{2}} \sin^{N-2} \theta \beta_0(\cos \theta) d\theta \mathbf{1}_{\frac{|z|}{\varepsilon} \leq 1} \\ &\propto \frac{1}{\varepsilon^N} J\left(\frac{|z|}{\varepsilon}\right) \end{aligned}$$

Notice that  $\int S_\varepsilon(|z|) dz = \frac{1}{\varepsilon^N} \int J\left(\frac{|z|}{\varepsilon}\right) dz$  does not depend on  $\varepsilon$ .  
 $\text{Supp}(S_\varepsilon) \subset \mathcal{B}(0, \varepsilon)$

## Computation of $S$

$$S(|z|) = \lambda\delta_0 + S_1(|z|)$$

**Proof.**

$B_\varepsilon := \frac{\beta_0}{|z|^N} \mathbf{1}_{|z| \geq \varepsilon}$  gives  $S_\varepsilon$  which goes to  $S_0$  when  $\varepsilon \rightarrow 0$

$$\begin{aligned} S_\varepsilon(|z|) &\propto \frac{1}{|z|^N} \int_0^{\frac{\pi}{2}} \sin^{N-2} \theta \left( \mathbf{1}_{|z| \geq \varepsilon \cos \frac{\theta}{2}} - \mathbf{1}_{|z| \geq \varepsilon} \right) \beta_0(\cos \theta) d\theta \\ &\propto \frac{1}{\varepsilon^N} \frac{1}{(|z|/\varepsilon)^N} \int_{2 \cos^{-1} \frac{|z|}{\varepsilon}}^{\frac{\pi}{2}} \sin^{N-2} \theta \beta_0(\cos \theta) d\theta \mathbf{1}_{\frac{|z|}{\varepsilon} \leq 1} \\ &\propto \frac{1}{\varepsilon^N} J\left(\frac{|z|}{\varepsilon}\right) \end{aligned}$$

Notice that  $\int S_\varepsilon(|z|) dz = \frac{1}{\varepsilon^N} \int J\left(\frac{|z|}{\varepsilon}\right) dz$  does not depend on  $\varepsilon$ .  
 $\text{Supp}(S_\varepsilon) \subset B(0, \varepsilon)$   
We get  $S_0 \propto \delta_0$

## Idea of the proof of strong compacity appearance

1. get strong compacity results in  $L^1_{(\text{loc})}$  for  $\int \beta(f^n)\phi(v)dv$ ,  
 $\beta(f^n) \star_v \phi$  and  $\sqrt{f^n} \star_v \phi$ .  $\implies$  AVERAGING LEMMAS  
strong convergence:  $\int f^n dv \rightarrow \int f dv$

## Idea of the proof of strong compacity appearance

1. get strong compacity results in  $L^1_{(\text{loc})}$  for  $\int \beta(f^n) \phi(v) dv$ ,  
 $\beta(f^n) \star_v \phi$  and  $\sqrt{f^n} \star_v \phi$ .  $\implies$  AVERAGING LEMMAS  
strong convergence:  $\int f^n dv \rightarrow \int f dv$
2. apply result of previous paper (*truncation*) and OVERALL ANGULAR SINGULARITY CONDITION  $\implies$  *velocity smoothness*:  $\lim_{A \rightarrow \infty} \overline{\lim}_{n \rightarrow \infty} \int_{W^\varepsilon} \int_{|\xi| \geq A} |\mathcal{F}\sqrt{f_R^n}|^2 d\xi dt dx$  where  
 $\sqrt{f^n} \rightarrow F$  and  $W_\varepsilon = \{(t, x) \in [0; T] \times B_R^N, \int F^2 dv > \varepsilon\}$

## Idea of the proof of strong compacity appearance

1. get strong compacity results in  $L^1_{(\text{loc})}$  for  $\int \beta(f^n) \phi(v) dv$ ,  
 $\beta(f^n) \star_v \phi$  and  $\sqrt{f^n} \star_v \phi$ .  $\implies$  AVERAGING LEMMAS  
strong convergence:  $\int f^n dv \rightarrow \int f dv$
2. apply result of previous paper (*truncation*) and OVERALL ANGULAR SINGULARITY CONDITION  $\implies$  velocity smoothness:  
 $\lim_{A \rightarrow \infty} \overline{\lim}_{n \rightarrow \infty} \int_{W^\varepsilon} \int_{|\xi| \geq A} |\mathcal{F}\sqrt{f_R^n}|^2 d\xi dt dx$  where  
 $\sqrt{f^n} \rightarrow F$  and  $W_\varepsilon = \{(t, x) \in [0; T] \times B_R^N, \int F^2 dv > \varepsilon\}$
3. use mollifiers to get strong compacity of  $(\sqrt{f^n})$  in  $L^2$

## Idea of the proof of strong compacity appearance

1. get strong compacity results in  $L^1_{(\text{loc})}$  for  $\int \beta(f^n) \phi(v) dv$ ,  
 $\beta(f^n) \star_v \phi$  and  $\sqrt{f^n} \star_v \phi$ .  $\implies$  AVERAGING LEMMAS  
strong convergence:  $\int f^n dv \rightarrow \int f dv$
2. apply result of previous paper (*truncation*) and OVERALL ANGULAR SINGULARITY CONDITION  $\implies$  velocity smoothness:  
 $\lim_{A \rightarrow \infty} \overline{\lim_{n \rightarrow \infty}} \int_{W^\varepsilon} \int_{|\xi| \geq A} |\mathcal{F}\sqrt{f_R^n}|^2 d\xi dt dx$  where  
 $\sqrt{f^n} \rightarrow F$  and  $W_\varepsilon = \{(t, x) \in [0; T] \times B_R^N, \int F^2 dv > \varepsilon\}$
3. use mollifiers to get strong compacity of  $(\sqrt{f^n})$  in  $L^2$
4. get  $L^1$  strong compacity of  $(f^n)$  as in DiPerna&Lions

## Idea of the proof of strong compacity appearance

1. get strong compacity results in  $L^1_{(\text{loc})}$  for  $\int \beta(f^n) \phi(v) dv$ ,  
 $\beta(f^n) \star_v \phi$  and  $\sqrt{f^n} \star_v \phi$ .  $\implies$  AVERAGING LEMMAS  
strong convergence:  $\int f^n dv \rightarrow \int f dv$
2. apply result of previous paper (*truncation*) and OVERALL ANGULAR SINGULARITY CONDITION  $\implies$  velocity smoothness:  
 $\lim_{A \rightarrow \infty} \overline{\lim}_{n \rightarrow \infty} \int_{W^\varepsilon} \int_{|\xi| \geq A} |\mathcal{F}\sqrt{f_R^n}|^2 d\xi dt dx$  where  
 $\sqrt{f^n} \rightarrow F$  and  $W_\varepsilon = \{(t, x) \in [0; T] \times B_R^N, \int F^2 dv > \varepsilon\}$
3. use mollifiers to get strong compacity of  $(\sqrt{f^n})$  in  $L^2$
4. get  $L^1$  strong compacity of  $(f^n)$  as in DiPerna&Lions
5. pass to the (strong) limit in the renormalized formulation,  
using the previous decomposition, and  $\mathcal{T}^n \rightarrow \mathcal{T}$  and *Fatou's lemma*

## Idea of the proof of strong compacity appearance

1. get strong compacity results in  $L^1_{(\text{loc})}$  for  $\int \beta(f^n) \phi(v) dv$ ,  
 $\beta(f^n) \star_v \phi$  and  $\sqrt{f^n} \star_v \phi$ .  $\implies$  AVERAGING LEMMAS  
strong convergence:  $\int f^n dv \rightarrow \int f dv$
2. apply result of previous paper (*truncation*) and OVERALL ANGULAR SINGULARITY CONDITION  $\implies$  velocity smoothness:  
 $\lim_{A \rightarrow \infty} \lim_{n \rightarrow \infty} \int_{W^\varepsilon} \int_{|\xi| \geq A} |\mathcal{F}\sqrt{f_R^n}|^2 d\xi dt dx$  where  
 $\sqrt{f^n} \rightarrow F$  and  $W_\varepsilon = \{(t, x) \in [0; T] \times B_R^N, \int F^2 dv > \varepsilon\}$
3. use mollifiers to get strong compacity of  $(\sqrt{f^n})$  in  $L^2$
4. get  $L^1$  strong compacity of  $(f^n)$  as in DiPerna&Lions
5. pass to the (strong) limit in the renormalized formulation,  
using the previous decomposition, and  $\mathcal{T}^n \rightarrow \mathcal{T}$  and *Fatou's lemma*
6. Conclude: check that  $f$  is a RSDM

# *Non-isotropic norm and estimates of linearized collision operator*

Norm used to get existence of *classical solutions*

# Non-isotropic norm and estimates of linearized collision operator

Norm used to get existence of *classical solutions*

$$\mu(v) = (2\pi)^{-\frac{3}{2}} e^{-\frac{|v|^2}{2}}$$

$$f = \mu + \sqrt{\mu}g$$

$$Q(f, f) = Q(\mu, \sqrt{\mu}g) +$$

$$Q(\sqrt{\mu}g, \mu) + Q(\sqrt{\mu}, \sqrt{\mu})$$

$$\Omega(g, h) := \frac{1}{\sqrt{\mu}} Q(\sqrt{\mu}g, \sqrt{\mu}h)$$

$$\mathcal{L}g := -\Omega(\sqrt{\mu}, g) - \Omega(g, \sqrt{\mu})$$

$$\text{linear: } \text{Ker}\mathcal{L} = \mathcal{N}$$

$\implies \mathbf{P}$   $L^2$ -orthogonal projection  
on  $\mathcal{N}$

Boltzmann equation  $\implies$

$$\partial_t g + v \cdot \nabla_x g + \mathcal{L}g = \Omega(g, g)$$

# Non-isotropic norm and estimates of linearized collision operator

Norm used to get existence of *classical solutions*

$$\mu(v) = (2\pi)^{-\frac{3}{2}} e^{-\frac{|v|^2}{2}}$$

$$f = \mu + \sqrt{\mu}g$$

$$Q(f, f) = Q(\mu, \sqrt{\mu}g) +$$

$$Q(\sqrt{\mu}g, \mu) + Q(\sqrt{\mu}, \sqrt{\mu})$$

$$\Omega(g, h) := \frac{1}{\sqrt{\mu}} Q(\sqrt{\mu}g, \sqrt{\mu}h)$$

$$\mathcal{L}g := -\Omega(\sqrt{\mu}, g) - \Omega(g, \sqrt{\mu})$$

$$\text{linear: } \text{Ker}\mathcal{L} = \mathcal{N}$$

$\implies$  **P**  $L^2$ -orthogonal projection  
on  $\mathcal{N}$

Boltzmann equation  $\implies$

$$\partial_t g + v \cdot \nabla_x g + \mathcal{L}g = \Omega(g, g)$$

## Non-isotropic norm

$$\begin{aligned} |||g|||^2 &= \int \Phi b \mu_*(g' - g)^2 dv_* dv d\sigma + \int \Phi b g_*^2 (\sqrt{\mu'} - \sqrt{\mu})^2 dv_* dv d\sigma \\ &= J_1 + J_2 \end{aligned}$$

# Non-isotropic norm and estimates of linearized collision operator

Norm used to get existence of *classical solutions*

$$\mu(v) = (2\pi)^{-\frac{3}{2}} e^{-\frac{|v|^2}{2}}$$

$$f = \mu + \sqrt{\mu}g$$

$$Q(f, f) = Q(\mu, \sqrt{\mu}g) +$$

$$Q(\sqrt{\mu}g, \mu) + Q(\sqrt{\mu}, \sqrt{\mu})$$

$$\Omega(g, h) := \frac{1}{\sqrt{\mu}} Q(\sqrt{\mu}g, \sqrt{\mu}h)$$

$$\mathcal{L}g := -\Omega(\sqrt{\mu}, g) - \Omega(g, \sqrt{\mu})$$

$$\text{linear: } \text{Ker}\mathcal{L} = \mathcal{N}$$

$\implies \mathbf{P}$   $L^2$ -orthogonal projection  
on  $\mathcal{N}$

Boltzmann equation  $\implies$

$$\partial_t g + v \cdot \nabla_x g + \mathcal{L}g = \Omega(g, g)$$

## Non-isotropic norm

$$\begin{aligned} |||g|||^2 &= \int \Phi b \mu_*(g' - g)^2 dv_* dv d\sigma + \int \Phi b g_*^2 (\sqrt{\mu'} - \sqrt{\mu})^2 dv_* dv d\sigma \\ &= J_1 + J_2 \end{aligned}$$

## Theorem

$B(z, \sigma) = \Phi(|z|)b(\cos \theta)$  with  $\Phi(|z|) = |z|^\gamma$  and similar

assumptions as before:  $\gamma > -3$  and  $\nu \in ]0; 2[$ .

$$C_1 |||(\mathbf{I} - \mathbf{P})g|||^2 \leq (\mathcal{L}g, g)_{L^2(\mathbb{R}_v^3)} \leq C_2 |||g|||^2$$

## About the non-isotropic norm

$$C_3 \left( \|g\|_{H^{\frac{\nu}{2}}(\mathbb{R}^3_\nu)}^2 + \|g\|_{L^{\frac{\nu+\gamma}{2}}(\mathbb{R}^3_\nu)}^2 \right) \leq |||g|||^2 \leq C_4 \|g\|_{H^{\frac{\gamma}{2}}(\mathbb{R}^3_\nu)}^2$$

## About the non-isotropic norm

$$C_3 \left( \|g\|_{H^{\frac{\gamma}{2}}(\mathbb{R}^3_\nu)}^2 + \|g\|_{L^2_{\frac{\nu+\gamma}{2}}(\mathbb{R}^3_\nu)}^2 \right) \leq |||g|||^2 \leq C_4 \|g\|_{H^{\frac{\gamma}{2}}_{\frac{\nu+\gamma}{2}}(\mathbb{R}^3_\nu)}^2$$

Treat separately  $J_1$  and  $J_2$

Show in particular that  $J_2^\Phi \sim \|g\|_{L^2_{\frac{\nu+\gamma}{2}}}^2 \sim J_2^{\tilde{\Phi}}$  and

$$J_1^\Phi + \|g\|_{L^2_{\frac{\nu+\gamma}{2}}}^2 \sim J_1^{\tilde{\Phi}} + \|g\|_{L^2_{\frac{\nu+\gamma}{2}}}^2$$

## About the non-isotropic norm

$$C_3 \left( \|g\|_{H^{\frac{\nu}{2}}(\mathbb{R}^3_v)}^2 + \|g\|_{L^2_{\frac{\nu+\gamma}{2}}(\mathbb{R}^3_v)}^2 \right) \leq |||g|||^2 \leq C_4 \|g\|_{H^{\frac{\gamma}{2}}_{\frac{\nu+\gamma}{2}}(\mathbb{R}^3_v)}^2$$

Treat separately  $J_1$  and  $J_2$

Show in particular that  $J_2^\Phi \sim \|g\|_{L^2_{\frac{\nu+\gamma}{2}}}^2 \sim J_2^{\tilde{\Phi}}$  and

$$J_1^\Phi + \|g\|_{L^2_{\frac{\nu+\gamma}{2}}}^2 \sim J_1^{\tilde{\Phi}} + \|g\|_{L^2_{\frac{\nu+\gamma}{2}}}^2$$

$\tilde{\Phi}(|z|) = <|z|>^\gamma = (1+|z|^2)^{\frac{\gamma}{2}} \implies$  get rid off the singularity in 0.

# Conclusion

Thank you !

Any questions ?