

2000 Zurich
proseminar
guidance

The Luttinger Liquid Concept via Bosonization


Christoph Buchendorfer

Abstract

It is shown how the Hamiltonian of the 1-dimensional spinless Luttinger model can be reexpressed in terms of boson operators. Even when one allows for interactions, the resulting boson Hamiltonian will be that of a free boson field. An example is given that shows how bosonization can simplify calculations of correlation functions. The additional features appearing when one allows for spin such as spin-charge separation are stated.

1 The Luttinger Model

The Luttinger model describes electrons of a 1-dimensional system with the following linear dispersion relation:



The states of the lower triangle have been introduced for mathematical reasons as will be apparent later and do not correspond to physical states. Electrons on the right branch are called right movers and those on the left branch left movers. Notation:

1 THE LUTTINGER MODEL

- \oplus : right-movers $c_{\oplus,k}^\dagger$
- \ominus : left-movers $c_{\ominus,k}^\dagger$

These are the creation operators for the corresponding branches. In order to refer to right and left movers by a variable index the letter r will be used. A value of 1 corresponds to \oplus and -1 corresponds to \ominus .

The corresponding continuous functions are defined as:

$$\psi_{\oplus}^\dagger(x) := \frac{1}{\sqrt{L}} \sum_k c_{\oplus,k}^\dagger e^{ikx} \quad (1)$$

$$\psi_{\ominus}^\dagger(x) := \frac{1}{\sqrt{L}} \sum_k c_{\ominus,k}^\dagger e^{ikx} \quad (2)$$

The ground state $|\phi_0\rangle$ is characterized by

$$\langle n_{r,k} \rangle_0 = \Theta(k_F - rk) \quad (3)$$

Normal ordering with respect to ground state:

$$: A : \equiv A - \langle A \rangle_0 \quad (4)$$

The free Luttinger Hamiltonian is then given by

$$H_0 = v_F : \sum_{k,r} (rk - k_F) (c_{r,k}^\dagger c_{r,k}) : \quad (5)$$

Equivalently

$$H_0 = v_F \int_0^L dx : \psi_{\oplus}^\dagger(i\partial_x - k_F)\psi_{\oplus} - \psi_{\ominus}^\dagger(i\partial_x + k_F)\psi_{\ominus} : \quad (6)$$

We define the density operators:

$$\psi_{\oplus}^\dagger(x)\psi_{\oplus}(x) =: \rho_{\oplus}(x) =: \frac{1}{L} \sum_q \rho_{\oplus,q} e^{iqx} \quad (7)$$

$$\psi_{\ominus}^\dagger(x)\psi_{\ominus}(x) =: \rho_{\ominus}(x) =: \frac{1}{L} \sum_q \rho_{\ominus,q} e^{iqx} \quad (8)$$

The free Hamiltonian can also be expressed in terms of density operators:

$$H_0 = \frac{\pi v_F}{L} : \sum_q \{ \rho_{\oplus,q} \rho_{\oplus,-q} + \rho_{\ominus,-q} \rho_{\ominus,q} \} : \quad (9)$$

The interaction part in the Luttinger model is given by

$$H_I = \frac{1}{2L} : \sum_{q \neq 0} 2g_2 \rho_{\oplus,q} \rho_{\ominus,-q} + g_4 (\rho_{\oplus,q} \rho_{\oplus,-q} + \rho_{\ominus,-q} \rho_{\ominus,q}) : \quad (10)$$

g_2 and g_4 are free parameters of the model.

2 Link to a lattice model

By means of numerous approximations one can map a 1-dimensional Hubbard model with nearest neighbour interaction onto a Luttinger model.

- one-dimensional lattice of length L , M lattice sites,
- lattice point spacing s ,
- periodic boundary conditions,
- half filling
- spinless fermions
- Hubbard-Hamiltonian with nearest neighbour hopping in the kinetic part,
- repulsive fermion-fermion interaction

$$H = H_0 + H_I \quad (11)$$

where

$$H_0 := -\varepsilon \sum_j \psi_j^\dagger \psi_{j+1} + \psi_{j+1}^\dagger \psi_j \quad (12)$$

$$H_I := \frac{U}{2} \sum_j n_j n_{j+1}, \quad U > 0 \quad (13)$$

ψ_i^\dagger is a fermionic creation operator that creates a fermion in a state whose wavefunction is localized around lattice point i ; ψ_i is its corresponding annihilation operator. They satisfy the common anti-commutator relations:

$$\{\psi_i, \psi_j^\dagger\} = \delta_{ij} \quad (14)$$

$$\{\psi_i, \psi_j\} = 0 \quad (15)$$

One can think of the wavefunctions as *Wannier-functions*, which are defined as superpositions of the exact (non interacting fermions) Bloch-wavefunctions:

$$\Phi_k = e^{ikx} u_k(x), \quad H_0 \Phi_k = \varepsilon(k) \Phi_k \quad \text{Bloch} \quad (16)$$

$$\omega(x - js) := \frac{1}{M} \sum_k e^{-ik(js)} \Phi_k(x) \quad \text{Wannier} \quad (17)$$

Where k is of the form

$$k = \frac{2\pi}{L} n, \quad n \in \mathbb{N} \quad (18)$$

Wannier-functions are localized around their corresponding lattice site.

2 LINK TO A LATTICE MODEL

4

In terms of the Fourier coefficients c_k of ψ defined by

$$\psi_j = \frac{1}{\sqrt{M}} \sum_k c_k e^{ikj s} \quad (19)$$

the kinetic Hamiltonian H_0 yields:

$$H_0 = -2\varepsilon \sum_k \cos(ks) c_k^\dagger c_k \quad (20)$$

Thus the Energy spectrum of the noninteracting fermions is given by

$$\varepsilon(k) = -2\varepsilon \cos(ks) \quad (21)$$

We assume half filling. Then the Fermimomentum k_F yields

$$k_F = \frac{M\pi}{2L} = \frac{\pi}{2s} \quad (22)$$

Which lies half way from the 1. Brillouin zone Boundary.

For the fermi-velocity v_F we obtain:

$$v_F = \left. \frac{\partial \varepsilon(k)}{\partial k} \right|_{k=k_F} = 2\varepsilon s \sin(k_F s) = 2\varepsilon s \quad (23)$$

In order to proceed we will perform a continuum limit based on the assumption:

$$\frac{s}{L} \rightarrow 0 \quad (24)$$

- lattice spacing s is assumed to stay *finite*.
- continuum limit considers
 - macroscopic L
 - atomic s
- compared to L the lattice points ns can be regarded as values of a continuous variable x .

We are interested in the behaviour near k_F and linearize the energy spectrum in the continuum limit around $\pm k_F$:

$+k_F$

$$\begin{aligned} \varepsilon(k) &= \varepsilon(k_F) + 2\varepsilon s \sin(k_F s)(k - k_F) + O(s^3(k - k_F)^3) \\ &\approx v_F(k - k_F) \end{aligned}$$

$-k_F$

$$\begin{aligned}\varepsilon(k) &= \varepsilon(-k_F) + 2s \sin(-k_F s)(k + k_F) + O(s^3(k + k_F)^3) \\ &\approx v_F(-k - k_F)\end{aligned}$$

We work now with two major simplifications:

- Linear spectrum $v_F(\pm k - k_F)$ extended over the whole range of k .
- Addition of "unphysical" states with a spectrum that is

- linear

- unbounded below

These approximations restrict the validity of the model to

- **LOW TEMPERATURES**

We have arrived at the assumptions of the free Luttinger Hamiltonian.

The mapping of the interaction part is not shown here.

3 Bosonization

We first consider the free case without interaction.

In terms of the Fourier-components $c_{r,k}^\dagger, c_{r,k}$ of the fermi field the density operator is:

$$\rho_{r,q} = \begin{cases} \sum_k c_{r,k+q}^\dagger c_{r,k} & (q \neq 0) \\ : \sum_k n_{r,k} : & (q = 0) \end{cases} \quad (25)$$

The hermitian conjugate is given by:

$$\rho_{r,q}^\dagger = \rho_{r,-q} \quad (26)$$

as shows the following computation (r omitted)

$$\begin{aligned}\rho_p^\dagger &= \left(\sum_k c_{k+p}^\dagger c_k \right)^\dagger \\ &= \sum_k c_k^\dagger c_{k+p} \\ &= \sum_k c_{k-p}^\dagger c_k \\ &= \rho_{-p}\end{aligned}$$

A straightforward calculation gives the following commutator

$$[\rho_{r,q}, \rho_{r',-q'}] = -\delta_{r,r'} \delta_{q,q'} r \frac{Lq}{2\pi} \quad (27)$$

However this result that can only be obtained if the unphysical states are present:

Let's assume $q > 0$. Routine commutator manipulation leads to

$$[\rho_{\oplus,q}, \rho_{\oplus,-q'}] = \delta_{q,q'} \sum_k \langle n_{\oplus,k} \rangle_0 - \langle n_{\oplus,k-q} \rangle_0$$

If the number of occupied states were finite a shift in the summation index would turn the sum into 0.

Let us assume that there is a $k_0 < k_F$ with all levels below occupied. Above k_0 arbitrarily many particle-hole states may be excited:

$$\begin{aligned} \sum_k \langle n_{\oplus,k} \rangle_0 - \langle n_{\oplus,k-q} \rangle_0 &= - \left(\sum_{k \geq k_0} + \sum_{k < k_0} \right) (\langle n_{\oplus,k-q} \rangle_0 - \langle n_{\oplus,k} \rangle_0) \\ &= - \sum_{k \geq k_0} (\langle n_{\oplus,k-q} \rangle_0 - \langle n_{\oplus,k} \rangle_0) \\ &= - \left(\sum_{k \geq k_0-q} \langle n_{\oplus,k} \rangle_0 - \sum_{k \geq k_0} \langle n_{\oplus,k} \rangle_0 \right) \\ &= - \sum_{k_0-q \leq k < k_0} \langle n_{\oplus,k} \rangle_0 \\ &= - \frac{L}{2\pi} q \end{aligned}$$

The following operators turn out to be bosonic:

$$b_q^\dagger := \sqrt{\frac{2\pi}{L|q|}} \sum_r \Theta(rq) \rho_{r,q} \quad (28)$$

$$b_q := \sqrt{\frac{2\pi}{L|q|}} \sum_r \Theta(rp) \rho_{r,-q} \quad (29)$$

Reversively:

$$\rho_{r,q} = \delta_{q,0} n_r + \sqrt{\frac{L|q|}{2\pi}} \{ \Theta(rq) b_q^\dagger + \Theta(-rq) b_{-q} \} \quad (30)$$

The bosonic commutation relations are satisfied:

$$[b_q, b_{q'}^\dagger] = \delta_{q,q'} \quad [b_q^\dagger, b_{q'}^\dagger] = 0 \quad (31)$$

Proof:

$$\begin{aligned} [b_q, b_{q'}^\dagger] &= \frac{2\pi}{L} \frac{1}{\sqrt{|q||q'|}} \sum_{r,r'} \Theta(rq)\Theta(r'q') [\rho_{r,-q}, \rho_{r',q'}] \\ &= \delta_{q,q'} \sum_r \frac{rq}{|q|} \Theta(rq) \\ &= \delta_{q,q'} \end{aligned}$$

$$\begin{aligned} [b_q^\dagger, b_{q'}^\dagger] &= \frac{2\pi}{L} \frac{1}{\sqrt{|q||q'|}} \sum_{r,r'} \Theta(r,q)\Theta(r',q') [\rho_{r,q}, \rho_{r',q'}] \\ &= \frac{-1}{|q|} \delta_{q,-q'} \sum_r rq \Theta(rq)\Theta(-rq) \\ &= 0 \end{aligned}$$

- b_q annihilates the ground state

$$\begin{aligned} b_q |\phi_0\rangle &= \sqrt{\frac{2\pi}{L|q|}} \sum_r \Theta(rq) \rho_{r,-q} |\phi_0\rangle \\ &= \sqrt{(*)} \sum_{r,k} \Theta(rq) c_{r,k-q}^\dagger c_{r,k} |\phi_0\rangle \\ &\stackrel{q>0}{=} \sqrt{(*)} \sum_k c_{\oplus,k-q}^\dagger c_{\oplus,k} |\phi_0\rangle = 0 \\ &\stackrel{q<0}{=} \sqrt{(*)} \sum_k c_{\ominus,k+|q|}^\dagger c_{\ominus,k} |\phi_0\rangle = 0 \end{aligned}$$

- b_q^\dagger creates excited states but does not raise the fermion number.

The commutator $[b_q, H_0]$ is given by

$$[b_q, H_0] = v_F |q| b_q \quad (32)$$

The commutator of the corresponding construction operator is now simply

$$[H_0, b_q^\dagger] = [b_q, H_0]^\dagger = |q| v_F b_q^\dagger \quad (33)$$

We thus have the algebra of harmonic oscillators with $\omega(q) = |q| v_F$ and consequently the free Hamiltonian H_0 on the space \mathcal{H}_B spanned by the excited states with respect to b_q^\dagger is given by:

$$: H_{0B} := v_F \sum_q |q| b_q^\dagger b_q \quad (34)$$

It can be shown, that $\mathcal{H}_B = \mathcal{H}$, so that the boson states form a complete set. Then $H_0 \equiv H_{0B}$ and the fermionic system can completely be represented in terms of bosonic states.

So far we have not gained any advantages by using boson operators but merely got acquainted with them. This will change when we introduce the interaction of the model. $H = H_0 + H_I$ still corresponds to a free, massless bosonic field.

The following fields are defined:

$$\phi(x) := -i \frac{\pi}{L} \sum_q \frac{1}{q} e^{-\alpha \frac{|q|}{2}} e^{-iqx} (\rho_{\Theta,q} + \rho_{\ominus,q}) \quad (35)$$

$$\theta(x) := i \frac{\pi}{L} \sum_q \frac{1}{q} e^{-\alpha \frac{|q|}{2}} e^{-iqx} (\rho_{\Theta,q} - \rho_{\ominus,q}) \quad (36)$$

Commutator:

$$[\phi(x), \theta(y)] = i \frac{\pi}{2} \text{sign}(x - y) \quad (37)$$

The gradient of θ :

$$\nabla \theta = \frac{\pi}{L} \sum_q e^{-\alpha \frac{|q|}{2}} e^{-iqx} (\rho_{\Theta,q} - \rho_{\ominus,q}) \quad (38)$$

Define the field conjugate to ϕ :

$$\Pi := \frac{\nabla \theta}{\pi} \quad (39)$$

Commutator:

$$[\phi(x), \Pi(y)] = i \delta(x - y) \quad (40)$$

Define:

$$u := \frac{1}{2\pi} \sqrt{(2\pi v_F + g_4)^2 - g_2^2} \quad (41)$$

$$K := \sqrt{\frac{2\pi v_F + g_4 - g_2}{2\pi v_F + g_4 + g_2}} \quad (42)$$

The crucial steps in Bosonization are the following claims:

$$H =: \int_0^L dx \left\{ \frac{\pi u K}{2} \Pi^2 + \frac{u}{2\pi K} (\partial_x \phi)^2 \right\} : \quad (43)$$

$$\psi_r(x) \propto e^{-irk_F x} \frac{1}{\sqrt{2\pi\alpha}} e^{i[\theta - r\phi]} \quad (44)$$

The expression for the Hamiltonian can be verified by a straightforward calculation just inserting the expressions for the fields ϕ and Π

The Hamiltonian has the form of an elastic string, which in its standard form looks like:

$$H = \int dx \left\{ \frac{1}{2\mu} \Pi^2 + \frac{\varepsilon}{2} (\partial_x \phi)^2 \right\} \quad (45)$$

In our case

$$\mu = \frac{1}{\pi u K}, \quad \varepsilon = \frac{u}{\pi K} \quad (46)$$

The eigenfrequencies of such a string are given by

$$\omega_k = \sqrt{\frac{\varepsilon}{\mu}} |k|$$

which in our case evaluates to

$$\omega_k = u |k| = \frac{1}{2\pi} \sqrt{(2\pi v_F + g_4)^2 - g_2^2} |k| \quad (47)$$

We can immediately compute the *Specific Heat* out of that result:

With

$$U = \sum_q \varepsilon(k) \frac{1}{e^{\beta \varepsilon(k)} - 1}, \quad \varepsilon(k) = u |k|, \quad \beta := \frac{1}{T} \quad (48)$$

$$\begin{aligned} C_V &= \frac{\partial U}{\partial T} = \frac{1}{T^2} \sum_q \varepsilon^2(k) \frac{e^{\beta \varepsilon(k)}}{(e^{\beta \varepsilon(k)} - 1)^2} \\ &= \frac{u^2}{4T^2} \sum_q \frac{q^2}{\sinh^2\left(\frac{\beta u q}{2}\right)} \\ &\approx \frac{u^2}{4T^2} \frac{L}{2\pi} \left(\frac{2}{\beta u}\right)^3 \int_{-\infty}^{\infty} dx \frac{x^2}{\sinh^2(x)} \\ &= \frac{L\pi}{3} \frac{1}{u} T \end{aligned}$$

$$C_V = \frac{L\pi}{3} \frac{1}{u} T \quad (49)$$

For free fermions u must be replaced by v_F and we get

$$\frac{C_V}{C_{V0}} = \frac{v_F}{u} \quad (50)$$

Since in our model $g_4 = g_2$ and $g_2 \ll v_F$ we can approximate:

$$\frac{C_V}{C_{V0}} \approx 1 - \frac{g_2}{2\pi v_F} \quad (51)$$

Since here $g_2 > 0$ the specific heat decreases.

4 Free Massless Field

This section states some facts about free massless boson fields that we will need later on.

The Hamiltonian of the classical free massless field is given by

$$H_B = \frac{1}{2} \int_0^L dx (\Pi^2(x) + (\nabla\phi(x))^2) \quad (52)$$

Quantization is performed by replacing Π and ϕ by corresponding hermitian field-operators with their commutator being

$$[\phi(x), \Pi(y)] = i\delta(x - y) \quad (53)$$

The fields can be expressed in terms of the bosonic Operators $\phi(q)$, $\phi^\dagger(q)$:

$$\phi(x) = \int_{-\infty}^{\infty} dq \frac{1}{2\pi\sqrt{2|q|}} [\phi(q)e^{iqx} + \phi^\dagger(q)e^{-iqx}] \quad (54)$$

$$\Pi(x) = -i \int_{-\infty}^{\infty} dq \frac{|q|}{2\pi\sqrt{2|q|}} [\phi(q)e^{iqx} - \phi^\dagger(q)e^{-iqx}] \quad (55)$$

Here

$$[\phi(q), \phi^\dagger(p)] = 2\pi\delta(q - p) \quad (56)$$

The Hamiltonian now becomes

$$H_B = \int_{-\infty}^{\infty} dq \frac{1}{2\pi} |q| \phi^\dagger(q)\phi(q) \quad (57)$$

Later on we will need the following correlation function:

$$G(x) := \langle \phi(x)\phi(0) - \phi^2(0) \rangle = \frac{1}{4\pi} \ln \frac{\alpha^2}{\alpha^2 + x^2} \quad (58)$$

$$G(x) = \int_{-\infty}^{\infty} dq \frac{1}{2\pi\sqrt{2|q|}} \int_{-\infty}^{\infty} dp \frac{1}{2\pi\sqrt{2|p|}} \langle \phi(q)\phi^\dagger(p) \rangle (e^{iqx} - 1) e^{-\alpha\frac{|q|}{2}} e^{-\alpha\frac{|p|}{2}}$$

$$\begin{aligned}
&= \int_{-\infty}^{\infty} dq \frac{1}{4\pi |q|} (e^{iqx} - 1) \\
&= \frac{1}{4\pi} \int_0^{\infty} dq \frac{1}{q} (e^{iqx} - 1) \\
&\quad + \frac{1}{4\pi} \int_0^{\infty} dq \frac{1}{q} (e^{-iqx} - 1) \\
&= \frac{1}{4\pi} \left(\ln \frac{\alpha}{\alpha - ix} + \ln \frac{\alpha}{\alpha + ix} \right) \\
&= \frac{1}{4\pi} \ln \frac{\alpha^2}{\alpha^2 + x^2}
\end{aligned}$$

The following identity is important:

$$e^A e^B =: e^{A+B} : e^{\langle AB + \frac{A^2 + B^2}{2} \rangle} \quad (59)$$

This leads to

$$G_\beta := \langle e^{i\beta\phi(x)} e^{-i\beta\phi(0)} \rangle = \left(\frac{\alpha^2}{\alpha^2 + x^2} \right)^{\frac{\beta^2}{4\pi}} \quad (60)$$

$$\begin{aligned}
G_\beta &= \langle : e^{i\beta(\phi(x) - \phi(0))} : \rangle \\
&= e^{\beta^2 \langle \phi(x)\phi(0) - (\phi^2(0) + \phi^2(x))/2 \rangle} \\
&= e^{\frac{\beta^2}{4\pi} \ln \frac{\alpha^2}{\alpha^2 + x^2}} \\
&= \left(\frac{\alpha^2}{\alpha^2 + x^2} \right)^{\frac{\beta^2}{4\pi}}
\end{aligned}$$

Note that due to translational invariance

$$\langle \phi^2(0) \rangle = \langle \phi^2(x) \rangle$$

Renormalizing the field avoids vanishing of the correlator as α tends to zero:

$$[e^{i\beta\phi}]_R := \alpha^{-\frac{\beta^2}{4\pi}} e^{i\beta\phi} \quad (61)$$

$$G_{\beta,R} = \left(\frac{1}{\alpha^2 + x^2} \right)^{\frac{\beta^2}{4\pi}} \quad (62)$$

5 Correlator in Luttinger Model

The aim is to calculate the correlation function

$$\chi(x, t) := \langle \rho(x, t) \rho(0, 0) \rangle \quad (63)$$

$$\rho(x) := \rho_{\oplus} + \rho_{\ominus} + (\psi_{\oplus}^{\dagger} \psi_{\ominus} + \text{h.c.}) \quad (64)$$

$$\lim_{\xi \rightarrow 0} \psi_{\oplus}^{\dagger}(x + \xi) \psi_{\ominus}(x - \xi) = \frac{1}{4\pi\alpha} e^{i2k_F x} e^{i2\phi} \quad (65)$$

This follows immediately from the representations of the fermion-field in terms of the boson fields ϕ and θ , if the following commutator vanishes:

$$\lim_{\xi \rightarrow 0} [\phi(x + \xi) - \theta(x + \xi), \theta(x - \xi) + \phi(x - \xi)] = 0 \quad (66)$$

$$\begin{aligned} \text{L.H.S} &= [\phi(x + \xi), \theta(x - \xi)] + [\phi(x - \xi), \theta(x + \xi)] \\ &= i\frac{\pi}{2}(\text{sign}(x + \xi - x + \xi) + \text{sign}(x - \xi - x - \xi)) \\ &= i\frac{\pi}{2}(\text{sign}(2\xi) + \text{sign}(-2\xi)) \\ &= 0 \end{aligned}$$

Writing out the terms of χ we get the following types:

- $\langle \nabla\phi(x) \nabla\phi(0) \rangle$
- $\langle e^{i2\phi} \nabla\phi \rangle$
- $\langle e^{i2\phi(x)} e^{i2\phi(0)} \rangle$
- $\cos(2k_F) \langle e^{i2\phi(x)} e^{-i2\phi(0)} \rangle$

All but the last term can be disposed of by subtle arguments. Thus we only look at the last term.

In order to apply the results of the last section we have to work with

$$\begin{aligned} \tilde{\Pi} &:= \sqrt{\pi K} \Pi \\ \tilde{\phi} &:= \frac{1}{\pi\sqrt{K}} \phi \end{aligned}$$

In other words $\beta = 2\sqrt{\pi K}$. We get

$$\langle e^{i2\phi(x)} e^{-i2\phi(0)} \rangle = \left(\frac{\alpha^2}{\alpha^2 + x^2} \right)^K \quad (67)$$

For large x α in the denominator is neglectable. We obtain a power law:

$$\propto \left(\frac{1}{x} \right)^{2K} \quad (68)$$

6 Spin

If we introduce spin, every operator in the starting equations acquires a spin index. We now have

$$H_0 = v_F \sum_{k,r,s} (rk - k_F) (c_{r,k,s}^\dagger c_{r,k,s} - \langle n_{r,k,s} \rangle_0) \quad (69)$$

Backward scattering:

$$H_{I,1} = \frac{1}{L} \sum_{kpqst} g_1 c_{\oplus,k,s}^\dagger c_{\ominus,p,t}^\dagger c_{\oplus,p+2k_F+q,t} c_{\ominus,k-2k_F-q,s} \quad (70)$$

$$H_{I,2} = \frac{1}{2L} \sum_{qst} \{ 2g_2 \rho_{\oplus,q,s} \rho_{\ominus,-q,t} + g_4 [\rho_{\oplus,q,s} \rho_{\oplus,-q,t} + \rho_{\ominus,-q,s} \rho_{\ominus,q,t}] \} \quad (71)$$

One now defines fields ϕ and Π for the two spin projections \uparrow and \downarrow in analogy to the spinless case.

Out of these charge and spin bosons are constructed:

- charge bosons

$$\phi_\rho := \frac{\phi_\uparrow + \phi_\downarrow}{\sqrt{2}} \quad (72)$$

$$\Pi_\rho := \frac{\Pi_\uparrow + \Pi_\downarrow}{\sqrt{2}} \quad (73)$$

- spin bosons

$$\phi_\sigma := \frac{\phi_\uparrow - \phi_\downarrow}{\sqrt{2}} \quad (74)$$

$$\Pi_\sigma := \frac{\Pi_\uparrow - \Pi_\downarrow}{\sqrt{2}} \quad (75)$$

These obey:

$$[\phi_\nu(x), \Pi_\mu(y)] = i\delta_{\nu\mu}\delta(x-y) \quad (76)$$

The Hamiltonian becomes:

$$H = H_\rho + H_\sigma + \frac{2g_1}{(2\pi\alpha)^2} \int dx \cos(\sqrt{8}\phi_\sigma) \quad (77)$$

with

$$H_\nu = \int dx \left[\frac{\pi u_\nu K_\nu}{2} \Pi_\nu^2 + \frac{u_\nu}{2\pi K_\nu} (\partial_x \phi_\nu)^2 \right] \quad (78)$$

$$u_\nu := \sqrt{\left(v_F + \frac{g_{4,\nu}}{\pi}\right)^2 - \left(\frac{g_\nu}{2\pi}\right)^2}$$

$$K_\nu := \sqrt{\frac{2\pi v_F + 2g_{4,\nu} + g_\nu}{2\pi v_F + 2g_{4,\nu} - g_\nu}}$$

$$g_\rho := g_1 - 2g_2$$

$$g_\sigma := g_1$$

$$g_{4,\rho} := g_4$$

$$g_{4,\sigma} := 0$$

For a noninteracting system we have

$$u_\nu = v_F \quad (79)$$

that is charge and spin velocities are equal.

For $g_1 = 0$:

- charge and spin oscillations are independent

If $g_1 \neq 0$ one has to treat the cosine term perturbatively.

This gives rise to the phenomenon of

- spin charge separation

from the Heisenberg equation of motion one obtains due to the linear spectrum:

$$\rho(x,t) = \rho(x - u_\rho t)$$

$$\sigma(x,t) = \sigma(x - u_\sigma t)$$

Let us now consider the case of adding a particle at x_0 to the system in ground state:

$$\begin{aligned}\langle \phi_0 | \psi_{\oplus}(x_0) \rho(x) \psi_{\oplus}^{\dagger}(x_0) | \phi_0 \rangle &= \delta(x - x_0) \\ \langle \phi_0 | \psi_{\oplus}(x_0) \sigma_z(x) \psi_{\oplus}^{\dagger}(x_0) | \phi_0 \rangle &= \delta(x - x_0)\end{aligned}$$

Then at time t :

$$\begin{aligned}\langle \phi_0 | \psi_{\oplus}(x_0) \rho(x - u_{\rho}t) \psi_{\oplus}^{\dagger}(x_0) | \phi_0 \rangle &= \delta(x - x_0 - u_{\rho}t) \\ \langle \phi_0 | \psi_{\oplus}(x_0) \sigma_z(x - u_{\sigma}t) \psi_{\oplus}^{\dagger}(x_0) | \phi_0 \rangle &= \delta(x - x_0 - u_{\sigma}t)\end{aligned}$$

Since in the interaction case $u_{\rho} \neq u_{\sigma} = v_F$

- spin waves are unaffected by the interaction
- charge waves are affected
- charge and spin will separate completely

References

- F.D.M. Haldane, *Luttinger liquid theory of one-dimensional quantum fluids*, J.Phys. C: Solid State Phys., 14 (1989)2585-2609
- V.J. Emery *Highly Conducting One-Dimensional Solids*, Chapter 6, Plenum Press New York (1979)
- R. Shankar, *Bosonization: How to make it work for you in Condensed Matter*
- Handwritten summaries of K. Le Hur, ITP ETH-Zuerich