

## Retarding subdiffusion and accelerating superdiffusion governed by distributed-order fractional diffusion equations

A. V. Chechkin,<sup>1,\*</sup> R. Gorenflo,<sup>2,†</sup> and I. M. Sokolov<sup>3,4,‡</sup>

<sup>1</sup>*Institute for Theoretical Physics, National Science Center, Kharkov Institute of Physics and Technology, Akademicheskaya Street 1, 61108 Kharkov, Ukraine*

<sup>2</sup>*Erstes Mathematisches Institut, FB Mathematik & Informatik, Freie Universität Berlin, Arnimallee 3, D-14195 Berlin, Germany*

<sup>3</sup>*Institut für Physik, Humboldt-Universität zu Berlin, Invalidenstrasse 110, D-10115 Berlin, Germany*

<sup>4</sup>*Theoretische Polymerphysik, Universität Freiburg, Hermann-Herder-Strasse 3, D-79104 Freiburg im Breisgau, Germany*

(Received 7 June 2002; published 22 October 2002)

We propose diffusionlike equations with time and space fractional derivatives of the distributed order for the kinetic description of anomalous diffusion and relaxation phenomena, whose diffusion exponent varies with time and which, correspondingly, cannot be viewed as self-affine random processes possessing a unique Hurst exponent. We prove the positivity of the solutions of the proposed equations and establish their relation to the continuous-time random walk theory. We show that the *distributed-order time fractional diffusion equation* describes the subdiffusion random process that is subordinated to the Wiener process and whose diffusion exponent decreases in time (*retarding subdiffusion*). This process may lead to *superslow diffusion*, with the mean square displacement growing logarithmically in time. We also demonstrate that the distributed-order space fractional diffusion equation describes superdiffusion phenomena with the diffusion exponent increasing in time (*accelerating superdiffusion*).

DOI: 10.1103/PhysRevE.66.046129

PACS number(s): 02.50.Ey, 05.40.-a

### I. INTRODUCTION

Recently, kinetic equations with fractional space and time derivatives have attracted attention as a possible tool for the description of anomalous diffusion and relaxation phenomena, see, e.g., Refs. [1–4] and references on earlier studies therein. It was also recognized [5–9] that the fractional kinetic equations may be viewed as “hydrodynamic” (that is, long-time and long-space) limit of continuous-time random walks (CTRW) [10], a model that was successfully applied to describe anomalous diffusion phenomena in many areas, e.g., turbulence [11], disordered media [12], intermittent chaotic systems [13]. Kinetic equations have two advantages over a random walk approach: first, they allow one to explore various boundary conditions (e.g., reflecting and/or absorbing) and, second, to study diffusion and/or relaxation phenomena in external fields. Both possibilities are difficult to realize in the framework of CTRW.

There are three types of fractional kinetic equations: the first one, describing Markovian processes, contains equations with fractional space or velocity derivative; the second one, describing non-Markovian processes, contains equations with fractional time derivative; and the third class, naturally, contains both fractional space and time derivatives, as well. However, all three types are suitable to describe time evolution of the probability density function (PDF) of a very narrow class of diffusion processes, which are characterized by a unique diffusion exponent showing time dependence of the

characteristic displacement (e.g., of the root mean square) [1]. These processes are also called fractal, or self-affine processes, and are characterized by the exponent  $H$ , called the Hurst exponent, which depends on the order of fractional derivative in the kinetic equation. We recall that the stochastic process  $x(t)$  is self-affine, or fractal, if its stationary increments possess the following property [14]:

$$x(t + \kappa\tau) - x(t) = \kappa^H [x(t + \tau) - x(t)], \quad (1)$$

where  $\kappa$  and  $H$  are positive constants. The sign  $\stackrel{d}{=}$  implies that the left and the right hand sides of Eq. (1) have the same PDFs.

As a possible generalization of fractional kinetics, we propose fractional diffusion equations in which the fractional order derivatives are integrated with respect to the order of differentiation (*distributed-order fractional diffusion equations*). They can serve as a paradigm for the kinetic description of the random processes possessing a nonunique diffusion exponent and hence, a nonunique Hurst exponent. The processes with a time-dependent Hurst exponent are believed to provide useful models for a host of continuous and nonstationary natural signals; they are also constructed explicitly [15–17]. Ordinary differential equations with distributed-order derivatives were proposed in the works by Caputo [18,19] for generalizing the stress-strain relation of inelastic media. In Refs. [20], [21], the method of the solution based on the generalized Taylor series representation was proposed. A basic framework for the numerical solution of distributed-order differential equations was introduced in Ref. [22]. Very recently, Caputo [23] proposed the generalization of the Fick’s law using the distributed-order time derivative.

\*Electronic address: [achechkin@kipt.kharkov.ua](mailto:achechkin@kipt.kharkov.ua)

†Electronic address: [gorenflo@math.fu-berlin.de](mailto:gorenflo@math.fu-berlin.de)

‡Electronic address: [igor.sokolov@physik.hu-berlin.de](mailto:igor.sokolov@physik.hu-berlin.de)

**II. DISTRIBUTED-ORDER TIME FRACTIONAL DIFFUSION EQUATION**

We write the distributed order time fractional diffusion equation for the PDF  $f(x,t)$  as

$$\int_0^1 d\beta \tau^{\beta-1} p(\beta) \frac{\partial^\beta f}{\partial t^\beta} = D \frac{\partial^2 f}{\partial x^2}, \quad f(x,0) = \delta(x), \quad (2)$$

where  $\tau$  and  $D$  are positive constants,  $\tau$  is given in seconds and  $D$  is given in units of  $\text{cm}^2/\text{sec}$ ,  $p(\beta)$  is a dimensionless non-negative function, and the time fractional derivative of order  $\beta$  is understood in the Caputo sense [24],

$$\frac{\partial^\beta f}{\partial t^\beta} = \frac{1}{\Gamma(1-\beta)} \int_0^t dt' (t-t')^{-\beta} \frac{\partial f}{\partial t'}. \quad (3)$$

If we set  $p(\beta) = \delta(\beta - \beta_0)$ ,  $0 < \beta_0 \leq 1$ , we arrive at time fractional diffusion equation, whose solution is the PDF of the self-affine random process with the Hurst exponent equal to  $\beta_0/2$ . The PDF is expressed through the Wright function [25]. The diffusion process is then characterized by the mean square displacement

$$\langle x^2(t) \rangle \equiv \int_{-\infty}^{\infty} dx x^2 f(x,t) = \frac{2}{\Gamma(\beta_0+1)} D \tau^{1-\beta_0} t^{\beta_0}. \quad (4)$$

This formula provides the generalization of the corresponding formula for classical diffusion valid at  $\beta_0=1$ . For  $\beta < 1$ , Eq. (4) describes the process of slow diffusion, or subdiffusion.

Let us now prove that the solution of Eq. (2) is a PDF. The derivation here parallels the method used in Ref. [26], see also Ref. [27]. Its aim is to show that the random process whose PDF obeys Eq. (2) is subordinated to the Wiener process. Returning to Eq. (2) and applying the Laplace and Fourier transforms in succession,

$$\hat{f}(k,s) = \int_{-\infty}^{\infty} dx e^{ikx} \int_0^t dt e^{-st} f(x,t), \quad (5)$$

we get from Eq. (2)

$$\hat{f}(k,s) = \frac{1}{s} \frac{I(s\tau)}{I(s\tau) + k^2 D \tau}, \quad (6)$$

where

$$I(s\tau) = \int_0^1 d\beta (s\tau)^\beta p(\beta). \quad (7)$$

We note that under the conditions described above the function  $I(s\tau)$  is completely monotonic on the positive real axis, i.e., it is positive and the signs of its derivatives alternate. We rewrite Eq. (6) as follows:

$$\hat{f}(k,s) = \frac{1}{s} \int_0^\infty du e^{-u[I+k^2 D \tau]} = \int_0^\infty du e^{-uk^2 D \tau} \tilde{G}(u,s), \quad (8)$$

where

$$\tilde{G}(u,s) = \frac{I(s\tau)}{s} e^{-uI(s\tau)} \quad (9)$$

is the Laplace transform of a function  $G(u,t)$  whose properties will be specified below. Now,  $f(x,t)$  can be written as

$$\begin{aligned} f(x,t) &= \int_{-\infty}^{\infty} \frac{dk}{2\pi} e^{-ikx} \int_{Br} \frac{ds}{2\pi i} e^{st} \int_0^\infty du e^{-uk^2 D \tau} \tilde{G}(u,s) \\ &= \int_0^\infty du \frac{e^{-x^2/4\pi D \tau}}{\sqrt{4\pi u D \tau}} G(u,t). \end{aligned} \quad (10)$$

The function  $G(u,t)$  is the PDF providing the subordination transformation, from time scale  $t$  to time scale  $u$ . Indeed, at first we note that  $G(u,t)$  is normalized with respect to  $u$  for any  $t$ . Using Eq. (9) we get

$$\int_0^\infty du G(u,t) = L_s^{-1} \int_0^\infty du \left[ \frac{I}{s} e^{-ul} \right] = L_s^{-1} \left[ \frac{1}{s} \right] = 1, \quad (11)$$

where  $L_s^{-1}$  is an inverse Laplace transformation. Now, to prove the positivity of  $G(u,t)$ , it is sufficient to show that its Laplace transform  $\tilde{G}(u,s)$  is completely monotonic on the positive real axis [28]. The last statement arises from the observation that  $\tilde{G}(u,s)$  is a product of two completely monotonic functions,  $I/s$  and  $\exp(-ul)$ . The monotonicity of the former is obvious, whereas the monotonicity of the latter is an elementary consequence of the criterion 2 in Ref. [28], Chap. XIII, Sec. 4. Thus, we may conclude that the solution of Eq. (2) is a PDF, and that the random process, whose PDF obeys a distributed-order time fractional diffusion equation, is subordinated to the Gaussian process using operational time.

**III. SUBDIFFUSION WITH RETARDATION AND SUPERSLOW DIFFUSION**

Let us concentrate on the behavior of the second moment of the PDF, that is, on the mean square displacement (MSD). Using Eq. (6), we get

$$\langle x^2 \rangle(t) = \left\{ - \frac{\partial^2 \hat{f}(k,t)}{\partial k^2} \right\} \Bigg|_{k=0} = 2D \tau L_s^{-1} \left\{ \frac{1}{s I(s\tau)} \right\}. \quad (12)$$

Consider two fractional exponents in Eq. (2), namely, let

$$p(\beta) = B_1 \delta(\beta - \beta_1) + B_2 \delta(\beta - \beta_2), \quad (13)$$

where  $0 < \beta_1 < \beta_2 \leq 1$ ,  $B_1 > 0$ ,  $B_2 > 0$ . Inserting Eq. (13) into Eq. (12) we get, denoting  $b_1 = B_1 \tau^{\beta_1}$ ,  $b_2 = B_2 \tau^{\beta_2}$ :

$$\begin{aligned} \langle x^2 \rangle(t) &= 2D\tau L_s^{-1} \left\{ \frac{1}{s(b_1 s^{\beta_1} + b_2 s^{\beta_2})} \right\} \\ &= \frac{2D\tau}{b_2} L_s^{-1} \left\{ \frac{s^{-\beta_1-1}}{\frac{b_1}{b_2} + s^{\beta_2-\beta_1}} \right\}. \end{aligned} \quad (14)$$

Recalling the Laplace transform of the generalized Mittag-Leffler function  $E_{\mu,\nu}(z)$ ,  $\mu > 0$ ,  $\nu > 0$ , which can be conveniently written as [24]

$$L_t \{ t^\nu E_{\mu,\nu}(-\lambda t^\mu) \} = \frac{s^{\mu-\nu}}{s^\mu + \lambda}, \quad \text{Re } s > x|\lambda|^{1/\mu}, \quad (15)$$

we get from Eq. (14),

$$\langle x^2 \rangle = \frac{2D\tau}{b_2} t^{\beta_2} E_{\beta_2-\beta_1, \beta_2+1} \left( -\frac{b_1}{b_2} t^{\beta_2-\beta_1} \right). \quad (16)$$

To get asymptotics at small  $t$ , we use an expansion, which is, in fact, the definition of  $E_{\mu,\nu}(z)$ , see Ref. [29], Chap. XVIII, Eq. (19),

$$E_{\mu,\nu}(z) = \sum_{n=0}^{\infty} \frac{z^n}{\Gamma(\mu n + \nu)}, \quad (17)$$

which yields, in the leading order for the MSD,

$$\langle x^2 \rangle \approx \frac{2D\tau}{B_2 \Gamma(\beta_2 + 1)} \left( \frac{t}{\tau} \right)^{\beta_2} \propto t^{\beta_2}. \quad (18)$$

For large  $t$  we use the following expansion valid on the real negative axis, see Ref. [29], Chap. XVIII, Eq. (21),

$$E_{\mu,\nu}(z) = - \sum_{n=1}^N \frac{z^{-n}}{\Gamma(-\mu n + \nu)} + O(|z|^{-1-N}), \quad |z| \rightarrow \infty, \quad (19)$$

which yields

$$\langle x^2 \rangle \approx \frac{2D\tau}{B_1 \Gamma(\beta_1 + 1)} \left( \frac{t}{\tau} \right)^{\beta_1} \propto t^{\beta_1}. \quad (20)$$

Since  $\beta_1 < \beta_2$ , we have the effect of *diffusion with retardation*. We also note that the kinetic equation with two fractional derivatives of different orders appears quite naturally when describing subdiffusive motion in velocity fields [30]. In this case the orders of derivatives are  $\beta$  and  $\beta - 1$ , so that the situation differs from the one discussed above.

Now we consider a simple particular case which, in some sense, is opposite to the cases considered above. Namely, we put

$$p(\beta) = 1, \quad 0 \leq \beta \leq 1. \quad (21)$$

Inserting Eq. (21) into Eq. (7) we get

$$I(s\tau) = \frac{s\tau - 1}{\ln(s\tau)}, \quad (22)$$

and, using then Eq. (12),

$$\langle x^2 \rangle = 2D\tau \left\{ \ln \frac{t}{\tau} + \gamma + e^{t/\tau} E_1 \left( \frac{t}{\tau} \right) \right\}, \quad (23)$$

where  $\gamma = 0.5772\dots$  is the Euler constant, and

$$E_1(z) = \int_z^\infty dy \frac{e^{-y}}{y} \quad (24)$$

is the exponential integral. Using now the expansions valid on the positive real axis, see Ref. [31], Eqs. (5.1.11) and (5.1.51), respectively,

$$E_1(z) = -\gamma - \ln z - \sum_{n=1}^{\infty} \frac{(-z)^n}{nn!}, \quad z \rightarrow 0 \quad (25)$$

and

$$E_1(z) \approx \frac{e^{-z}}{z} \sum_{n=0}^N (-1)^n \frac{z^n}{n!}, \quad z \rightarrow \infty, \quad (26)$$

we get, retaining the main terms of the asymptotics at small and large times, respectively,

$$\langle x^2 \rangle \approx \begin{cases} 2D\tau \frac{t}{\tau} \ln \frac{\tau}{t}, & t \rightarrow 0 \\ 2D\tau \ln \left( \frac{t}{\tau} \right), & t \rightarrow \infty. \end{cases} \quad (27)$$

Thus, at small times we have slightly anomalous superdiffusion, whereas at large times we have *superslow diffusion*.

The superslow diffusion, for which MSD grows logarithmically with time,  $\langle x^2(t) \rangle \propto \ln^2 t$ , was observed in the Sinai model [32], in aperiodic environments [33], and in an iterated map [34]. The superslow diffusion was also found numerically in an area preserving a parabolic map on a cylinder [35]. Up to now, it was unclear, whether this type of anomaly (“strong anomaly,” by terminology of Ref. [34]) can be described within the framework of fractional kinetics. The example presented above demonstrates that the distributed-order fractional kinetic equations can serve as a tool for the description of a strong diffusional anomaly.

The formula (27) can be generalized to the case

$$p(\beta) = \begin{cases} \frac{1}{\beta_2 - \beta_1}, & 0 \leq \beta_1 \leq \beta \leq \beta_2 \leq 1 \\ 0, & \text{otherwise,} \end{cases} \quad (28)$$

and  $\int_0^\infty d\beta p(\beta) = 1$ . Inserting Eq. (28) into Eq. (7) and then into Eq. (12), we get

$$\begin{aligned} \langle x^2 \rangle(t) &= \frac{2D\tau}{\beta_2 - \beta_1} L_s^{-1} \left\{ \frac{\ln(s\tau)}{s[(s\tau)^{\beta_2} - (s\tau)^{\beta_1}]} \right\} \\ &= - \frac{2D\tau}{\beta_2 - \beta_1} \left\{ \frac{d}{d\delta} L_s^{-1} \left[ \frac{s^{-\delta-\beta_1}}{s^{\beta_2-\beta_1} - 1} \right] \right\} \left( \frac{t}{\tau} \right) \Big|_{\delta=1}. \end{aligned} \quad (29)$$

Recalling Eq. (15), the MSD can be written as

$$\langle x^2 \rangle = \frac{2D\tau}{\beta_2 - \beta_1} \left( \frac{t}{\tau} \right)^{\beta_2} \left\{ \ln \left( \frac{\tau}{t} \right) E_{\beta_2 - \beta_1, \beta_2 + 1} \left( \left( \frac{t}{\tau} \right)^{\beta_2 - \beta_1} \right) - \left( \frac{d}{d\delta} E_{\beta_2 - \beta_1, \beta_2 + \delta} \left( \left( \frac{t}{\tau} \right)^{\beta_2 - \beta_1} \right) \right) \Big|_{\delta=1} \right\}. \quad (30)$$

Using Eq. (17), we get an expansion for  $\langle x^2 \rangle$  at small  $t$ ,

$$\langle x^2 \rangle = \frac{2D\tau}{\beta_2 - \beta_1} \left( \frac{t}{\tau} \right)^{\beta_2} \sum_{n=0}^{\infty} \left( \frac{t}{\tau} \right)^{n(\beta_2 - \beta_1)} \left\{ \ln \left( \frac{\tau}{t} \right) + \psi(1 + \beta_2 + n(\beta_2 - \beta_1)) \right\} \Gamma^{-1}[1 + \beta_2 + n(\beta_2 - \beta_1)], \quad (31)$$

where  $\psi(\nu) = d(\ln \Gamma(\nu))/d\nu$  is the  $\psi$  function. At large  $t$  we explore the asymptotics valid on the real positive axis, see Ref. [29], Chap. XVIII, Eq. (22),

$$E_{\mu, \nu}(z) = \frac{1}{\mu} z^{(1-\nu)/\mu} \exp(z^{1/\mu}) - \sum_{n=1}^N \frac{z^{-n}}{\Gamma(\nu - \mu n)} + O(|z|^{-1-N}). \quad (32)$$

Using Eq. (32), Eq. (30) takes on the form

$$\langle x^2 \rangle \approx \frac{2D\tau}{\beta_2 - \beta_1} \left( \frac{t}{\tau} \right)^{\beta_1} \sum_{n=0}^N \left( \frac{t}{\tau} \right)^{-n(\beta_2 - \beta_1)} \left\{ \ln \left( \frac{t}{\tau} \right) - \psi(1 + \beta_2 - n(\beta_2 - \beta_1)) \right\} \Gamma^{-1}[1 + \beta_2 - n(\beta_2 - \beta_1)]. \quad (33)$$

If we set  $\beta_1 = 0, \beta_2 = 1$  in Eqs. (31) and (33), then we arrive at the same expansions that are obtained by inserting Eqs. (25) and (26) into Eq. (23), respectively. In particular, the leading terms of the series (31) and (33) at  $\beta_1 = 0, \beta_2 = 1$  coincide with Eq. (27).

#### IV. RELATION TO THE CONTINUOUS-TIME RANDOM WALK THEORY

The fractional diffusion equations with a given order of fractional time derivative are closely connected to the CTRW processes with the power-law distribution of waiting times between the subsequent steps [1,8]. Now we establish the connection between the distributed-order time fractional diffusion equations and more general CTRW situations. Recall the basic formula of the CTRW in the Fourier-Laplace space [11],

$$\hat{f}(k, s) = \frac{1 - \bar{w}(s)}{s} \frac{1}{1 - \hat{\psi}(k, s)}, \quad (34)$$

where  $\bar{w}(s)$  is the Laplace transform of the waiting-time PDF  $w(t)$ , and  $\hat{\psi}(k, s)$  is the Fourier-Laplace transform of the joint PDF of jumps and waiting times  $\psi(\xi, t)$ . Assume

the decoupled joint PDF,  $\psi(\xi, t) = \lambda(\xi)w(t)$ , and that the jump length variance is finite, that is, the Fourier transform of  $\lambda(\xi)$  is

$$\hat{\lambda}(k) \approx 1 - D\tau k^2 \quad (35)$$

to the lowest orders in  $k$ . Then, we consider the situations in which mean waiting time diverges, that is, at large  $t$  the waiting time PDF behaves as

$$w(t) \approx \tau^\beta / t^{1+\beta}, \quad 0 < \beta < 1, \quad (36)$$

and, consequently,

$$\bar{w}(s) \approx 1 - (s\tau)^\beta \quad (37)$$

at small  $s$ . If  $\beta$  is constant, then inserting Eqs. (37) and (35) into Eq. (34) and making an inverse Fourier-Laplace transform, we arrive at the time fractional diffusion equation. Now let us consider the case when  $\beta$  fluctuates. Indeed, for example, in the model called the Arrhenius cascade, which is inspired from studies of disordered systems, the unique  $\beta$  appears only under the assumption that the random trapping time is related to the random height of the well by the Arrhenius law [36]. In a more realistic model, this law gives only the average value of the trapping time. Thus, we may speculate that in order to take into account the fluctuations of the trapping time, we can introduce the conditional PDF

$$w(t|\beta) \approx \tau^\beta / t^{1+\beta}, \quad (38)$$

and the PDF  $p(\beta)$ , as well. Now, we have the relation

$$w(t) = \int_0^1 d\beta p(\beta) w(t|\beta), \quad (39)$$

where  $[0;1]$  is the whole interval for variations of  $\beta$ . We note that all waiting-time distributions with  $\beta \geq 1$  correspond to a similar behavior described by the first-order derivative. Then, for the  $\bar{w}(s)$  we have, instead of Eq. (37),

$$\bar{w}(s) \approx 1 - \int_0^1 d\beta (s\tau)^\beta p(\beta), \quad p(\beta) \geq 0, \quad \int_0^1 d\beta p(\beta) = 1. \quad (40)$$

Inserting Eqs. (40) and (35) into Eq. (34) we arrive at Eqs. (6) and (7). Thus, we see that the weight function  $p(\beta)$  has the meaning of the PDF.

The model with fluctuating  $\beta$  is, of course, only one of the possible interpretations of Eq. (39): the nonexact power-law behavior of the waiting-time PDF can physically have very different reasons. In particular, the representation (39) allows us to consider *regularly varying* waiting-time PDFs, i.e., those which behave as  $w(t) \propto t^{-1-\beta} g(t)$  at  $t \rightarrow \infty$ , where  $g(t)$  is a slowly varying function, e.g., any power of  $\ln t$  [28]. We are also able to consider waiting-time PDFs  $w(t)$  that show an approximately scaling behavior with the exponents *changing with time*. For such distributions the effective PDFs  $p(\beta)$  can be determined, and thus such nonperfectly scaling CTRWs can be described through distributed-order diffusion

equations. The formal inversion of Eq. (39) can follow by noting that  $tw(t)$  taken as a function of  $\ln t$  is the Laplace transform of the function

$$\phi(\beta) = \begin{cases} \tau^\beta p(\beta), & 0 \leq \beta \leq 1 \\ 0, & 1 < \beta < \infty. \end{cases}$$

Indeed,

$$\begin{aligned} tw(t) &= \int_0^\infty d\beta \phi(\beta) t^{-\beta} = \int_0^\infty d\beta \phi(\beta) \exp(-\beta \ln t) \\ &= L_\beta\{\phi(\beta)\}(\ln t). \end{aligned}$$

The function  $\phi(\beta)$  is thus given by

$$\phi(\beta) = L_u^{-1}\{e^u w(e^u)\}.$$

The value of  $\tau$  can then be found through the normalization condition  $\int_0^\infty \phi(\beta) \tau^{-\beta} = 1$ , which then defines the function  $p(\beta)$ . The description of the process through the distributed-order diffusion equation is possible whenever this function is non-negative and concentrated on  $0 \leq \beta \leq 1$ .

### V. DISTRIBUTED-ORDER SPACE FRACTIONAL DIFFUSION EQUATION

Now we turn to another type of fractional equation, namely, *distributed-order space fractional diffusion equation* which, in dimensional variables, takes on the form

$$\frac{\partial f}{\partial t} = \int_{0^+}^2 d\alpha D(\alpha) \frac{\partial^\alpha f}{\partial |x|^\alpha}, \quad f(x,0) = \delta(x), \quad (41)$$

where  $D$  is a (dimensional) function of the order of the derivative  $\alpha$ , and the Riesz space fractional derivative  $\partial^\alpha/\partial|x|^\alpha$  is understood through its Fourier transform  $\hat{\Phi}$  as

$$\hat{\Phi} \left( \frac{\partial^\alpha f}{\partial |x|^\alpha} \right) = -|k|^\alpha \hat{f}. \quad (42)$$

If we set  $D(\alpha) = K_{\alpha_0} \delta(\alpha - \alpha_0)$ , then we arrive at the space fractional diffusion equation, whose solution is a Lévy stable PDF of the self-affine stable process whose Hurst exponent is equal to  $1/\alpha_0$ . The PDF is expressed in terms of the Fox's  $H$  function [37,38]. In the general case  $D(\alpha)$  can be represented as

$$D(\alpha) = l^{\alpha-2} DA(\alpha), \quad (43)$$

where  $l$  (in centimeters) and  $D$  (in  $\text{cm}^2/\text{sec}$ ) are dimensional positive constants,  $A$  is a dimensionless non-negative function of  $\alpha$ . The equation that follows for the characteristic function from Eq. (41) has the solution

$$\hat{f}(k,t) = \exp \left\{ -\frac{Dt}{l^2} \int_0^2 d\alpha A(\alpha) (|k|l)^\alpha \right\}. \quad (44)$$

Note that the normalization condition

$$\int_{-\infty}^\infty dx f(x,t) = \hat{f}(k=0,t) = 1 \quad (45)$$

is fulfilled.

Consider the simple particular case

$$A(\alpha) = A_1 \delta(\alpha - \alpha_1) + A_2 \delta(\alpha - \alpha_2), \quad (46)$$

where  $0 < \alpha_1 < \alpha_2 \leq 2$ ,  $A_1 > 0$ ,  $A_2 > 0$ . Inserting Eq. (46) into Eq. (44) we have

$$\hat{f}(k,t) = \exp\{-a_1 |k|^{\alpha_1} t - a_2 |k|^{\alpha_2} t\}, \quad (47)$$

where  $a_1 = A_1 D/l^{2-\alpha_1}$ ,  $a_2 = A_2 D/l^{2-\alpha_2}$ . The characteristic function (47) is the product of two characteristic functions of the Lévy stable PDFs with the Lévy indexes  $\alpha_1$ ,  $\alpha_2$ , and the scale parameters  $a_1^{1/\alpha_1}$  and  $a_2^{1/\alpha_2}$ , respectively. Therefore, the inverse Fourier transformation of Eq. (47) gives the PDF which is the convolution of the two stable PDFs,

$$\begin{aligned} f(x,t) &= a_1^{-1/\alpha_1} a_2^{-1/\alpha_2} t^{-1/\alpha_1 - 1/\alpha_2} \\ &\times \int_{-\infty}^\infty dx' L_{\alpha_1,0} \left( \frac{x-x'}{(a_1 t)^{1/\alpha_1}} \right) L_{\alpha_2,0} \left( \frac{x'}{(a_2 t)^{1/\alpha_2}} \right), \end{aligned} \quad (48)$$

where  $L_\alpha(x)$  is the PDF of the symmetric Lévy stable law possessing the characteristic function

$$\hat{L}_{\alpha,0}(k) = \exp(-|k|^\alpha). \quad (49)$$

The PDF given by Eq. (48) is, obviously, positive, as the convolution of two positive PDFs. The PDF will be also positive, if the function  $A(\alpha)$  is represented as a sum of  $N$   $\delta$  functions multiplied by positive constants,  $N$  is a positive integer. Moreover, if  $A(\alpha)$  is a continuous positive function, then discretizing the integral in Eq. (41) by a Riemann sum and passing to the limit we can also conclude on the positivity of the PDF.

### VI. SUPERDIFFUSION WITH ACCELERATION

Since the mean square displacement diverges for the Lévy stable process, the anomalous superdiffusion can be characterized by the typical displacement  $\delta x$  of the diffusing particle [39],

$$\delta x \propto \langle |x|^q \rangle^{1/q}, \quad (50)$$

where  $\langle |x|^q \rangle$  is the  $q$ th absolute moment of the PDF obeying Eq. (41). For the stable process with the Lévy index  $\alpha$

$$\langle |x|^q \rangle = \begin{cases} C(q;\alpha) t^{q/\alpha}, & 0 < q < \alpha < 2 \\ \infty, & q \geq \alpha, \end{cases} \quad (51)$$

where the coefficient

$$C(q; \alpha) = \frac{2}{\pi q} (K_{\alpha} t)^{q/\alpha} \sin\left(\frac{\pi q}{2}\right) \Gamma(1+q) \Gamma\left(1 - \frac{q}{\alpha}\right) \quad (52)$$

was obtained in Ref. [39]. To evaluate the  $q$ th moment for the case given by Eq. (46),  $q < \alpha_1$ , we use the following expression, see e.g., Ref. [40]:

$$\langle |x|^q \rangle = \frac{2}{\pi} \Gamma(1+q) \sin\left(\frac{\pi q}{2}\right) \int_0^{\infty} dk (1 - \text{Re} \hat{f}(k, t))^{-q-1}. \quad (53)$$

We insert Eq. (47) into Eq. (53) and expand in series either  $\exp(-a_1 |k|^{\alpha_1} t)$  or  $\exp(-a_2 |k|^{\alpha_2} t)$ , with subsequent integration over  $k$ . As the result, for the  $q$ th moment we have expansions valid at  $q < \alpha_1$  and for small and large times, respectively,

$$\begin{aligned} \langle |x|^q \rangle &= \frac{2}{\pi q} (a_2 t)^{q/\alpha_2} \sin\left(\frac{\pi q}{2}\right) \Gamma(1+q) \Gamma\left(1 - \frac{q}{\alpha_2}\right) \left\{ 1 + \frac{q}{\Gamma\left(1 - \frac{q}{\alpha_2}\right)} \right. \\ &\quad \left. \times \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{\alpha_2 n!} a_1^n a_2^{-n\alpha_1/\alpha_2} \Gamma\left(\frac{n\alpha_1 - q}{\alpha_2}\right) t^{n(1 - \alpha_1/\alpha_2)} \right\}, \quad t \rightarrow 0, \end{aligned} \quad (54)$$

$$\begin{aligned} \langle |x|^q \rangle &\approx \frac{2}{\pi q} (a_1 t)^{q/\alpha_1} \sin\left(\frac{\pi q}{2}\right) \Gamma(1+q) \Gamma\left(1 - \frac{q}{\alpha_1}\right) \left\{ 1 + \frac{1}{\Gamma\left(1 - \frac{q}{\alpha_1}\right)} \right. \\ &\quad \left. \times \sum_{n=1}^N \frac{(-1)^{n+1}}{\alpha_1 n!} a_2^n \alpha_1^{-n\alpha_2/\alpha_1} \Gamma\left(\frac{n\alpha_2 - q}{\alpha_1}\right) t^{-n(\alpha_2/\alpha_1 - 1)} \right\}, \quad t \rightarrow \infty. \end{aligned} \quad (55)$$

One can see, that at small times the characteristic displacement grows as  $t^{1/\alpha_2}$ , whereas at large times it grows as  $t^{1/\alpha_1}$ . Thus, we have *superdiffusion with acceleration*.

## VII. CONCLUSIONS

In summary, we believe that the distributed-order fractional diffusion equations can serve as a useful tool for the description of complicated diffusion processes, for which the diffusion exponent can change in the course of time. Further investigations are needed to establish the connection between proposed kinetics and multifractality. On the other hand, the development of numerical schemes for solving distributed-

order kinetic equations and for modeling sample paths of the random processes governed by these equations is also of importance.

## ACKNOWLEDGMENTS

The authors thank J. Klafter and F. Mainardi for fruitful discussions. This work was supported by the INTAS Project No 00-0847. A.V.C also acknowledges the hospitality of the Department of Mathematics and Computer Science of the Free University of Berlin and support from the DAAD. I.M.S. is grateful to the Fonds der Chemischen Industrie for partial financial support.

- [1] R. Metzler and J. Klafter, Phys. Rep. **339**, 1 (2000).
- [2] E. Lutz, Phys. Rev. Lett. **86**, 2208 (2001).
- [3] F. Mainardi, Yu. Luchko, and G. Pagnini, Fract. Calc. Appl. Anal. **4**, 153 (2001).
- [4] A. V. Chechkin and V. Yu. Gonchar, Zh. Eksp. Teor. Fiz. **118**, 730 (2000) [JETP **91**, 635 (2000)].
- [5] R. Hilfer and L. Anton, Phys. Rev. E **51**, R848 (1995).
- [6] A. Compte, Phys. Rev. E **53**, 4191 (1996).
- [7] F. Mainardi, M. Raberto, R. Gorenflo, and E. Scalas, Physica A **287**, 468 (2000).

- [8] E. Barkai, R. Metzler, and J. Klafter, Phys. Rev. E **61**, 132 (2000).
- [9] E. Barkai, Chem. Phys. (to be published), e-print cond-mat/0108024.
- [10] E. W. Montroll and G. H. Weiss, J. Math. Phys. **6**, 167 (1965).
- [11] J. Klafter, A. Blumen, and M. F. Schlesinger, Phys. Rev. A **35**, 3081 (1987).
- [12] J.-P. Bouchaud and A. Georges, Phys. Rep. **195**, 127 (1990).
- [13] G. Zumofen and J. Klafter, Phys. Rev. E **47**, 851 (1993).
- [14] G. Samorodnitsky and M. S. Taqqu, *Stable Non-Gaussian Pro-*

- cesses (Chapman & Hall, New York, 1994).
- [15] R.-F. Peltier and J. Lévy Véhel, INRIA Research Report No. 2645, 1995 (unpublished).
- [16] A. Ayache and J. Lévy Véhel, in *Fractals: Theory and Applications in Engineering*, edited by M. Dekking, J. Lévy Véhel, E. Lutton, and C. Triccot (Springer-Verlag, Berlin, 1999), p. 17.
- [17] A. Ayache and J. Lévy Véhel, *Statist. Infer. Stoch. Proc.* **1-2**, 7 (2000).
- [18] M. Caputo, *Elasticità Dissipazione* (Zanichelli Printer, Bologna, 1969).
- [19] M. Caputo, *Annals Univ. Ferrara-Sez. VII-Sc. Mat.* **XLI**, 73 (1995).
- [20] R. L. Bagley and P. J. Torvik, *Int. J. Appl. Math.* **2**, 865 (2000).
- [21] R. L. Bagley and P. J. Torvik, *Int. J. Appl. Math.* **2**, 965 (2000).
- [22] K. Diethelm and N. J. Ford, *Fract. Calc. Appl. Anal.* **4**, 531 (2001).
- [23] M. Caputo, *Fract. Calc. Appl. Anal.* **4**, 421 (2001).
- [24] R. Gorenflo and F. Mainardi, in *Fractals and Fractional Calculus in Continuum Mechanics*, edited by A. Carpinteri and F. Mainardi (Springer-Verlag, Vienna, 1997), p. 223.
- [25] F. Mainardi, in *Fractals and Fractional Calculus in Continuum Mechanics* (Ref. [24]), p. 291.
- [26] I. M. Sokolov, *Phys. Rev. E* **63**, 056111 (2001).
- [27] E. Barkai, *Phys. Rev. E* **63**, 046118 (2001).
- [28] W. Feller, *An Introduction to Probability Theory and Its Applications*, 2nd ed. (Wiley, New York, 1971), Vol. II.
- [29] A. Erdélyi, *Bateman Manuscript Project Higher Transcendental Functions* (McGraw-Hill, New York, 1955), Vol. III.
- [30] R. Metzler, J. Klafter, and I. M. Sokolov, *Phys. Rev. E* **58**, 1621 (1998).
- [31] *Handbook of Mathematical Functions*, edited by M. Abramowitz and I. A. Stegun (Dover, New York, 1965).
- [32] Y. G. Sinai, *Theor. Probab. Appl.* **27**, 256 (1982).
- [33] F. Igloi, L. Turban, and H. Rieger, *Phys. Rev. E* **59**, 1465 (1999).
- [34] J. Dräger and J. Klafter, *Phys. Rev. Lett.* **84**, 5998 (2000).
- [35] T. Prosen and M. Žnidarič, *Phys. Rev. Lett.* **87**, 114101 (2001).
- [36] F. Bardou, in *Proceedings of the Conference on Lévy Processes. Theory and Applications*, University of Aarhus, 1999, edited by O. E. Barndorff-Nielsen, S. E. Graversen, and T. Mikosch (unpublished), pp. 22–38 (downloadable from <http://www.maphysto.dk/publications/MPS-misc/1999/11.pdf>).
- [37] C. Fox, *Trans. Am. Math. Soc.* **98**, 395 (1961).
- [38] W. R. Schneider, in *Stochastic Processes in Classical and Quantum Systems*, edited by S. Albeverio, G. Casati, and D. Merlini, *Lecture Notes in Physics* (Springer-Verlag, Berlin, 1986), p. 497.
- [39] B. J. West and V. Seshadri, *Physica A* **113**, 203 (1982).
- [40] V. M. Zolotarev, *One-dimensional Stable Distributions* (American Mathematical Society, Providence, RI, 1986), Vol. 65.