

$$\text{IvP} \quad y' + \frac{2}{x}y = \frac{\cos x}{x^2}, \quad y(\pi) = 0$$

standard form of First order linear ODE is

$$\frac{dy}{dx} + P(x)y = Q(x), \quad y(x_0) = y_0$$

① clearly $P(x) = \frac{2}{x}$: $Q(x) = \frac{\cos x}{x^2}$ are continuous

for all $x \in (-\infty, 0) \cup (0, \infty)$.

Given that $x_0 = \pi$. ($\pi > 0$)

\therefore Consider IvP in $(0, \infty)$.

So, the IvP has unique solution in $(0, \infty)$.

② consider $\frac{dy}{dx} + \frac{2}{x}y = 0, \quad x \in (0, \infty)$

$$y' + \frac{2}{x}y = 0, \quad y > 0$$

$$\frac{y'}{y} = -\frac{2}{x}$$

on integrating from x_0 to x

$$\int_{x_0}^x \left(\frac{y'}{y} \right) dt = - \int_{x_0}^x \frac{2}{t} dt$$

$$\log y(t) \Big|_{x_0}^x = -2 \log t \Big|_{x_0}^x$$

$$\begin{aligned} \log y(x) - \log y(x_0) \\ = -2(\log x - \log x_0) \end{aligned}$$

$$\Rightarrow \log y(x) = -2 \log x + \log c$$

where

$$\log c = 2 \log x_0 + \log y(x_0).$$

$$\log y(x) = \log(c/x^2)$$

$$y(x) = \frac{c}{x^2}, \quad x \in (0, \infty),$$

$$c \in \mathbb{R} \setminus \{0, \infty\}.$$

Verification:

clearly $y(x) = \frac{c}{x^2}$ is differentiable on $(0, \infty)$.

$$y'(x) = \frac{-2c}{x^3}, \quad c \in \mathbb{R} \setminus \{0, \infty\}$$

$$\text{So, } y' + \frac{2}{x}y = -\frac{2c}{x^3} + \frac{2}{x} \times \frac{c}{x^2} = 0$$

$\therefore y(x) = \frac{c}{x^2}$ is a solution of $y' + \frac{2}{x}y = 0$.

③ Consider $\frac{dy}{dx} + \frac{2}{x}y = \frac{\cos x}{x^2}, \quad x \in (0, \infty)$

Let $y_p = \phi = \frac{c(x)}{x^2}$, where $c(x)$ is differentiable on $(0, \infty)$

$$\phi' = c'(x) + x^{-2} + c(x)x^{-2}x^{-3} \quad ; \quad y_p \text{ is a particular solution}$$

$$\text{So, } \phi' + \frac{2}{x}\phi = \frac{\cos x}{x^2}$$

$$\Rightarrow c'(x) \times x^2 - \frac{2}{x^3} f(x) + \frac{2}{x} \times \frac{f(x)}{x^2} = \frac{\cos x}{x^2}$$

$$\Rightarrow \frac{c'(x)}{x^2} = \frac{\cos x}{x^2}, \text{ since } x \neq 0.$$

$$\Rightarrow c'(x) = \cos x \Rightarrow c(x) = \sin x + D, \quad D \in \mathbb{R}.$$

\therefore Particular Solution is

$$y_p(x) = \phi(x) = \frac{1}{x^2} c(x) = \frac{1}{x^2} \left[\sin x + \frac{D}{x} \right]$$

$$= \frac{D}{x^2} + \frac{\sin x}{x^2}.$$

\therefore General solution of the non-homogeneous equation

$$\text{is } y = y_c + y_p$$

$$= \frac{C}{x^2} + \frac{D}{x^2} + \frac{\sin x}{x^2}$$

$$= \frac{c_1}{x^2} + \frac{\sin x}{x^2}, \text{ where } c_1 = (C+D) \in \mathbb{R}.$$

$$\text{Given } y(\pi) = 0.$$

$$\text{So, } 0 = \frac{c_1}{\pi^2} + \frac{\sin \pi}{\pi^2} \Rightarrow 0 = \frac{c_1}{\pi^2} + 0 \Rightarrow c_1 = 0$$

$$y = \frac{\sin x}{x^2}$$

(4) Verification: we have $y = \frac{c_1}{x^2} + \frac{\sin x}{x^2}, x \in (0, \infty)$

$$y' = -\frac{2c_1}{x^3} + \frac{2c_1}{x^3} + \frac{\cos x}{x^2} - \frac{2\sin x}{x^3}$$

Consider L.H.S

$$y' + \frac{2}{x} y = -\frac{2c_1}{x^3} + \frac{2c_1}{x^3} + \frac{\cos x}{x^2} - \frac{2\sin x}{x^3} + \frac{2\sin x}{x^2}$$

$$= \frac{\cos x}{x^2}. = \text{R.H.S.}$$

$\therefore y = \frac{c_1}{x^2} + \frac{\sin x}{x^2}$ is a ^{general} solution of

$$y' + \frac{2}{x} y = \frac{\sin x}{x^2} \frac{\cos x}{x^2}, x \in (0, \infty).$$