

Problem 5.

i) i). With Taylor's theorem up to the order 0 with the explicit formula for the remainder we have:

For every $\alpha \in [a, b], x \in [a, b]$, there is a $c \in (\alpha, x)$
 $x > \alpha$ such that $f(x) = f(\alpha) + f'(c) \cdot (x - \alpha)$

so if $f'(c) \geq 0$ for any $c \in \mathbb{R}$,

$$f(x) - f(\alpha) = \frac{f'(c)}{\geq 0} \cdot \frac{(x - \alpha)}{\geq 0}$$

$$\geq 0$$

so $f(x) \geq f(\alpha)$ when $x > \alpha$

ie f is non-decreasing on $[a, b]$

ii). With $f: [0, \frac{\pi}{2}] \rightarrow \mathbb{R}$

$$x \mapsto x - \sin(x)$$

f is of class C^1 , and $f'(x) = 1 - \cos(x)$ for any $x \in \mathbb{R}$

but $\forall t \in \mathbb{R}, -1 \leq \cos(t) \leq 1$, so $f'(x) \geq 0$

Hence f is non-decreasing on $[0, \frac{\pi}{2}]$.

$$\text{so } \forall x \in [0, \frac{\pi}{2}], f(x) \geq f(0) = 0 - \sin(0) = 0 - 0 = 0$$

Conclusion On $[0, \frac{\pi}{2}]$, $\sin(x) \leq x$

2)

Now, let's use the Taylor's theorem

up to the order 3 with the explicit formula for the remainder:

For $x \in (0, \pi/2)$, there is a $c \in (0, \pi)$ such that

$$\begin{aligned}\sin(x) &= \sin(0) + x \cdot \sin'(0) + \frac{x^2}{2} \cdot \sin''(0) \\ &\quad + \frac{x^3}{6} \cdot \sin'''(0) + \frac{x^4}{24} \cdot \sin^{(4)}(c)\end{aligned}$$

$\cdot \sin(0) = 0$

$\cdot \sin'(0) = \cos(0) = 1$

$\cdot \sin''(0) = -\sin(0) = 0$ and $\sin^{(4)}(0) = \sin$

$\cdot \sin'''(0) = -\cos(0) = -1$

So $\sin(\pi) = \pi - \frac{\pi^3}{6} + \frac{\pi^4}{24} \cdot \sin(c)$

But on $(0, \pi/2)$, $\sin(c) > 0$,

$\frac{\pi^4}{24} > 0$

so $\sin(\pi) = \pi - \frac{\pi^3}{6} + \frac{\pi^4}{24} \sin(c)$

$\therefore \pi - \frac{\pi^3}{6} \geq 0$

Conclusion

On $(0, \pi/2)$,

$$\pi - \frac{\pi^3}{6} \leq \sin(\pi).$$

3) These two inequalities

are still true for $x > \pi/2$.

* $\sin(\pi) \leq \pi$ is obvious if $\pi \geq 1$ because $\sin(\pi) \leq 1$

* Bonus: solve yourself the last inequality!