For these problems, you will study the curves with as much details as possible: domain of definition, quick drawing, asymptotes, special points, singular points, double points, extrema for x(t), y(t), $\sqrt{x^2(t) + y^2(t)}$ etc.

<u>Remark:</u> this tutorial sheet is the last one for the MA101 course.

Ex.1) Study these parametric curves (be cautious on the domain of t):

(a)
$$\begin{cases} x(t) = t - \sin(t) \\ y(t) = 1 - \cos(t) \end{cases}$$
(b)
$$\begin{cases} x(t) = \frac{4t}{3 + t^4} \\ y(t) = \frac{4}{\sqrt{3}} \frac{t^3}{3 + t^4} \end{cases}$$
(c)
$$\begin{cases} x(t) = \cos^3(t) \\ y(t) = \sin^3(t) \end{cases}$$
(d)
$$\begin{cases} x(t) = \frac{e^t}{1 + t} \\ y(t) = \frac{te^t}{1 + t} \end{cases}$$

If you are curious, you can try to look for these curves on Internet, and maybe find their names. A good website for plotting this is www.wolframalpha.com.

Ex.2) Study these polar curves (be cautious on the domain of θ):

(a)
$$\rho(\theta) = \cos(\theta) + \sin(2\theta)$$
,

(b)
$$\rho(\theta) = \cos(\theta) - \cos(2\theta)$$
,

(c)
$$\rho(\theta) = \frac{1 - 2\cos(\theta)}{1 + \sin(\theta)},$$

(d)
$$\rho(\theta) = \frac{\cos(2\theta)}{2\cos(\theta) - 1}$$

- (e) $\rho(\theta) = 1 + \tan\left(\frac{\theta}{2}\right)$, prove the existence of one *double point* on which the two tangents are orthogonal.
- (f) $\rho(\theta) = \frac{1}{\sin^3(\theta/3)}$, show that one line, passing to the origin O (ie the point $(0,0) \in \mathbb{R}^2$) will intersect the curve in exactly three points, and that the three tangents to the polar curve (at these points) will form a *equilateral triangle*.

Ex.3) Study the polar curve : $\rho(\theta) = \frac{1}{\sqrt{1 + \sin(2\theta)} + \sqrt{1 - \sin(2\theta)}}$. First, try to do it without cheating or plotting the curve on a calculator machine or a computer.

If any interesting behavior seems to appear, try to find a way to prove it^2 .

1

Please, report any issue to MA101@crans.org

¹Try it on Wolfram Alpha: www.wolframalpha.com/input/?i=plot(x(t)=1-sin(t),y(t)=1-cos(t))

²Here, one could try to manipulate trigonometric expressions to simplify the expression of ρ and then use Cartesian coordinates to get the final result.