

nekk

3c)

$$\frac{1}{1-3x+2x^2} = \frac{1}{(1-x)(1-2x)} = \frac{2}{1-2x} - \frac{1}{1-x}$$

$$= 2 \sum_{n=0}^{+\infty} (2x)^n - \sum_{n=0}^{+\infty} x^n$$

$|2x| < 1$        $|x| < 1$

pour  $x \in ]-\frac{1}{2}, \frac{1}{2}[$

$$= \sum_{n=0}^{+\infty} (2 \cdot 2^n - 1) x^n = \sum_{n=0}^{+\infty} (2^{n+1} - 1) x^n$$

Calcul du rayon :  $2^{-\frac{1}{2}} \cdot 2^{\frac{1}{2}} = R$ .

donc  $\sum_{n=0}^{+\infty} (2^{n+1} - 1) x^n$  et  $\sum_{n=0}^{+\infty} 2^{n+1} x^n$  ont le même rayon

$$\left| \frac{u_{n+1}}{u_n} \right| = \frac{2^{n+2}}{2^{n+1} |x|} = 2|x|$$

Si  $2|x| < 1$  alors  $\sum u_n$  converg.

$$R = \frac{1}{2}$$

$$\boxed{\frac{1}{1-3x+2x^2} = \sum_{n=0}^{+\infty} (2^{n+1} - 1) x^n, \forall x \in ]-\frac{1}{2}, \frac{1}{2}[}$$

$$\frac{1}{2-x} = \frac{1}{2(1-\frac{x}{2})} = \frac{1}{2} \sum_{n=0}^{+\infty} \left(\frac{x}{2}\right)^n = \sum_{n=0}^{+\infty} \frac{1}{2^{n+1}} x^n$$

pour  $x \in ]-1, 1[$ , donc pour  $x \in ]-2, 2[$

④  $4xy'' + 2y' + y = 0$

$$f(x) = \sum_{n=0}^{+\infty} a_n x^n$$

$$f'(x) = \sum_{n=1}^{+\infty} n a_n x^{n-1}$$

$$f''(x) = \sum_{n=1}^{+\infty} n(n-1) a_n x^{n-2}$$

$$4x f''(x) + 2f'(x) + f(x) = 4x \sum_{n=1}^{+\infty} n(n-1) a_n x^{n-2} + 2 \sum_{n=1}^{+\infty} n a_n x^{n-1} + \sum_{n=0}^{+\infty} a_n x^n = 0$$

$$= \sum_{n=1}^{+\infty} 4n(n-1) a_n x^{n-1} + \sum_{n=1}^{+\infty} 2na_n x^{n-1} + \sum_{n=0}^{+\infty} a_n x^n = 0$$

$$= \sum_{n=0}^{+\infty} (4(n+1)n a_{n+1} + 2(n+1)a_{n+1} + a_n) x^n = 0 = \sum_{n=0}^{+\infty} 0 x^n$$

$$\forall n \geq 0, 4(n+1)a_{n+1} + 2(n+1)a_{n+2} + a_n = 0$$

on décale de 1 :

$$\forall n \geq 1, 4n(n-1)a_n + 2na_n + a_{n-1} = 0$$

$$2n(2(n-1)a_n + a_{n-1}) + a_n = 0$$

$$\boxed{\forall n \geq 1, 2n(2n-1)a_n + a_{n-1} = 0}$$

$$n=1: 2 \cdot 1 \cdot a_1 + a_0 = 0 \Rightarrow a_1 = -\frac{a_0}{2}$$

$$n=2: 4 \cdot 3 \cdot a_2 + a_1 = 0 \Rightarrow a_2 = -\frac{a_1}{3 \cdot 4} = \frac{a_0}{2 \cdot 3 \cdot 4} = \frac{a_0}{4!}$$

$$n=3: 6 \cdot 5 \cdot a_3 + a_2 = 0 \Rightarrow a_3 = -\frac{a_2}{5 \cdot 6} = -\frac{a_0}{2 \cdot 3 \cdot 4 \cdot 5 \cdot 6} = -\frac{a_0}{6!}$$

Il semble que  $a_n = \frac{(-1)^n}{(2n)!} a_0$

Si  $a_0 = 1$ :  $f(x) = \sum_0^{+\infty} \frac{(-1)^n}{(2n)!} x^n$  =  $\sum_0^{+\infty} \frac{(-1)^n}{(2n)!} (\sqrt{x})^{2n}$  qd  $x \geq 0$  car  $(\sqrt{x})^2 = x$   
 $= \cos(\sqrt{x})$

$$\text{qd } x \leq 0: f(x) = \sum_0^{+\infty} \frac{(-x)^n}{(2n)!} = \sum_0^{+\infty} \frac{(\sqrt{-x})^{2n}}{(2n)!} = \underline{\text{ch}(\sqrt{x})}$$

ATS 2021 Exo 3 partie B

$$xy''(x) + y'(x) + \omega^2 x y(x) = 0 \quad \omega > 0$$

$$\text{en } \underline{x=0}: 0 + y'(0) + 0 = 0 \Rightarrow \underline{y'(0)=0}$$

$$y = \sum_0^{+\infty} a_n x^n \quad y' = \sum_1^{\infty} a_n n x^{n-1} = a_1 + 2a_2 x + 3a_3 x^2 + \dots$$

$$y'(0) = \underline{a_1 = 0}$$

$$y''(x) = \sum_1^{\infty} a_n n(n-1)x^{n-2}$$

$$xy'' + y' + \omega^2 x y = \sum_1^{\infty} a_n n(n-1)x^{n-1} + \sum_0^{\infty} n a_n x^{n-1} + \sum_0^{\infty} \omega^2 a_n x^{n+1}$$

$$= \sum_0^{\infty} a_{n+1}(n+1)n x^n + \sum_0^{\infty} (n+1)a_{n+1} x^n + \sum_1^{\infty} \omega^2 a_n x^n$$

$$= \sum_1^{\infty} a_{n+1}(n+1)n x^n + \sum_1^{\infty} (n+1)a_{n+1} x^n + \sum_1^{\infty} \omega^2 a_n x^n = 0$$

$$\sum_{n=1}^{\infty} \left[ \alpha_{n+1} (n+1) \omega + (n+1) \alpha_{n+1} + \omega^2 \alpha_{n-1} \right] x^n = 0$$

D'après l'unicité du développement en Séries Entières :

$$\forall n \geq 1 \quad (n+1)^2 \alpha_{n+1} + \omega^2 \alpha_{n-1} = 0 \Rightarrow \alpha_{n+1} = \frac{-\omega^2 \alpha_{n-1}}{(n+1)^2}$$

$$\Rightarrow \forall n \geq 0 \quad \alpha_{n+2} = \frac{-\omega^2 \alpha_n}{(n+2)^2}$$

$$\text{Montons par récurrence que } \alpha_{2n} = \frac{(-1)^n \omega^{2n}}{2^{2n} n!^2} \alpha_0$$

3c)  $\frac{1}{1-3x+2x^2} = \frac{1}{(1-x)(1-2x)} = \frac{-1}{1-x} + \frac{2}{1-2x}$

$= -\sum_{n=0}^{+\infty} x^n + 2 \sum_{n=0}^{+\infty} (Rx)^n$

$= -\sum_0^{\infty} x^n + 2 \sum_0^{\infty} 2^n x^n$

$= \sum_0^{+\infty} [2^{n+1} - 1] x^n$

$\forall x \in ]-\frac{1}{2}; \frac{1}{2}[ \quad R = \frac{1}{2}$

S. E da  $f(x) = \frac{1}{2-x} = \frac{1}{2(1-\frac{x}{2})} = \frac{1}{2} \sum_0^{+\infty} \left(\frac{x}{2}\right)^n \quad \forall x \in ]-2, 2[$

$= \sum_0^{+\infty} \frac{x^n}{2^{n+1}}$

$$\begin{aligned}
 \textcircled{1} \quad 4xy'' + 2y' + y &= 0 & f(x) &= \sum_{n=0}^{+\infty} a_n x^n & f'(x) &= \sum_{n=1}^{+\infty} n a_n x^{n-1} \\
 & & f''(x) &= \sum_{n=2}^{+\infty} n(n-1) a_n x^{n-2} & \\
 4xy'' + 2y' + y &= \sum_1^{\infty} 4n(n-1)a_n x^{n-1} + \sum_1^{\infty} 2na_n x^{n-1} + \sum_0^{+\infty} a_n x^n & & & \\
 & \downarrow & & \downarrow & \\
 &= \sum_0^{\infty} 4(n+1)n a_{n+1} x^n + \sum_0^{\infty} 2(n+1) a_{n+1} x^n + \sum_0^{+\infty} a_n x^n & & & \\
 &= \sum_0^{\infty} \left[ 4(n+1)n a_{n+1} + 2(n+1) a_{n+1} + a_n \right] x^n & & & \\
 &= \sum_0^{\infty} \left[ 2(n+1)(2n+1) a_{n+1} + a_n \right] x^n = 0 = \sum_0^{\infty} 0 x^n
 \end{aligned}$$

D'après l'uniformité des Dsrlt en S.E:  $\forall n \geq 0 \quad 2(n+1)(2n+1) a_{n+1} + a_n = 0$

on décale les n:

$$\text{pour } n=1: \quad 2a_2 + a_0 = 0$$

$$\text{pour } n=2: \quad 4 \cdot 3 a_3 + a_1 = 0$$

$$\text{pour } n=3: \quad 6 \cdot 5 a_4 + a_2 = 0$$

$$\boxed{\forall n \geq 1 \quad 2n(2n-1) a_n + a_{n-1} = 0}$$

$$a_0 = 1$$

$$a_1 = -\frac{1}{2} = \frac{(-1)^1}{2!}$$

$$a_2 = -\frac{a_1}{3 \cdot 4} = \frac{-1}{2 \cdot 3 \cdot 4} = \frac{(-1)^2}{4!}$$

$$a_3 = -\frac{a_2}{5 \cdot 6} = \frac{-1}{2 \cdot 3 \cdot 4 \cdot 5 \cdot 6} = \frac{(-1)^3}{6!}$$

Il semble que  $a_n = \frac{(-1)^n}{(2n)!}$  Recurrence :

Initialisation : pour  $n=0$

Hypothèse : Je prends un  $n \geq 0$ . Je suppose que  $a_n = \frac{(-1)^n}{(2n)!}$

$$a_{n+1} = \frac{-a_n}{(2n+2)(2n+1)} = \frac{-\frac{(-1)^n}{(2n)!}}{(2n+2)(2n+1)} = \frac{-(-1)^n}{(2n)! \cdot (2n+1)(2n+2)} = \frac{(-1)^{n+1}}{(2n+2)!}$$

Conclusion :  $\forall n \geq 0 \quad a_n = \frac{(-1)^n}{(2n)!}$

$$f(x) = \sum_{n=0}^{+\infty} \frac{(-1)^n}{(2n)!} x^{2n}$$

$$\text{Si } x \geq 0 : \cos(\sqrt{x}) = \sum_{n=0}^{+\infty} \frac{(-1)^n}{(2n)!} (\sqrt{x})^{2n} = \sum_{n=0}^{+\infty} \frac{(-1)^n}{(2n)!} x^n = f(x)$$

$$\text{Si } x \leq 0 : \operatorname{ch}(\sqrt{-x}) = \sum_{n=0}^{+\infty} \frac{1}{(2n)!} (\sqrt{-x})^{2n} = \sum_{n=0}^{+\infty} \frac{(-x)^n}{(2n)!} = \sum_{n=0}^{+\infty} \frac{(-1)^n x^n}{(2n)!} = f(x)$$

ARS 2021 exo 3 Route B

$$x y'' + y' + \omega^2 x y(x) = 0$$

① Montrer que  $y'(0) = 0$

$$\text{en } x=0 : 0 + y'(0) + 0 = 0 \Rightarrow \underline{y'(0)=0}$$

②  $y(x) = \sum_{n=0}^{+\infty} a_n x^n$ . Que vaut  $a_1$  ?

$$\text{On dérive : } y'(x) = \sum_{n=0, n \geq 1}^{+\infty} n a_n x^{n-1} = 0 + a_1 + 2a_2 x + 3a_3 x^2 + \dots$$

$$y'(0) = \underline{a_1} = 0$$

③ Montrer que  $\forall n \geq 0 \quad a_{n+2} = -\frac{\omega^2 a_n}{(n+2)^2}$

$$x y''(x) + y'(x) + \omega^2 x y(x) = \sum_{n=0, n \geq 2}^{+\infty} n(n-1) a_n x^{n-1} + \sum_{n=0}^{+\infty} n a_n x^{n-1} + \sum_{n=0}^{+\infty} \omega^2 a_n x^{n+1}$$

$$\sum_{n=1}^{+\infty} (n+1)n a_{n+1} x^n + \sum_{n=1}^{+\infty} (n+1) a_{n+1} x^n + \sum_{n=1}^{+\infty} \omega^2 a_{n+1} x^n = 0$$

$\downarrow$   
 $\downarrow$   
 $\downarrow a_1 = 0$

$$\sum_{n=1}^{+\infty} \left[ (n+1) \alpha_{n+1} + (n+1) \alpha_{n+1} + \omega^2 \alpha_{n-1} \right] x^n = 0 = \sum_{n=1}^{+\infty} 0 x^n$$

$\Rightarrow 0 \quad (\text{unicité des déréts en S. E.})$

$$(n+1) \alpha_{n+1} (n+1) + \omega^2 \alpha_{n-1} = 0$$

$$\forall n \geq 1: (n+1)^2 \alpha_{n+1} + \omega^2 \alpha_{n-1} = 0$$

on décale de 1

$$\alpha_{n+2} = \frac{-\omega^2 \alpha_n}{(n+2)^2}$$

$$\forall n \geq 0: (n+2)^2 \alpha_{n+2} + \omega^2 \alpha_n = 0 \quad \text{CQFD}$$

$$\text{M que } \alpha_{2n} = \frac{(-1)^n \omega^{2n}}{2^{2n} (n!)^2} \alpha_0 \text{ par récurrence}$$