

useee

$$\begin{array}{r}
 \textcircled{7} \quad \left| \begin{array}{r}
 2x^2 - 6x + 5 \\
 - (2x^2 - 2x) \\
 \hline
 -4x + 5 \\
 - (-4x + 4) \\
 \hline
 1
 \end{array} \right| \quad \begin{array}{l}
 x - 1 \\
 \hline
 2x - 4
 \end{array} \quad 2x^2 - 6x + 5 = (x-1)(2x-4) + 1
 \end{array}$$

~~$\frac{2x^2 - 6x + 5}{x-1} = 2x - 4 + 1$~~

$$\left(\frac{2x^2 - 6x + 5}{x-1} = 2x - 4 + \frac{1}{x-1} \right)$$

$$① a = \frac{1}{2 + \frac{1}{1 + \frac{1}{3+a}}}$$

$$\frac{3+a}{3+a} + \frac{1}{3+a} = \frac{4+a}{3+a}$$

$$a = \frac{1}{2 + \frac{1}{\frac{4+a}{3+a}}} = \frac{1}{2 + \frac{3+a}{4+a}} = \frac{1}{\frac{8+2a+3a}{4+a}} = \frac{4+a}{11+3a}$$

$$a = \frac{4+a}{11+3a} \quad (\Rightarrow a(11+3a) = 4+a \Rightarrow 3a^2 + 10a - 4 = 0)$$

$$(\Rightarrow 3a^2 + 10a - 4 = 0) \qquad \Delta = \dots$$

$$\textcircled{b} \quad 9 + \dots + 37 = \sum_{k=3}^{18} (2k+1) = 2 \left(\sum_{k=3}^{18} k \right) + \sum_{k=3}^{18} 1$$

(1^{er} + dernier) (nb de termes)

$$= 2 \cdot \frac{(18+3)(18-3+1)}{2} + 18 - 3 + 1 = \dots$$

\textcircled{c} $\binom{n}{p} = \frac{n!}{p!(n-p)!}$ Dans le triangle de Pascal, on additionne des entiers, donc c'est un entier

$$(2+\sqrt{5})^n = \binom{n}{0} 2^n + \binom{n}{1} 2^{n-1} \sqrt{5} + \binom{n}{2} 2^{n-2} (\sqrt{5})^2 + \dots + \binom{n}{n} (\sqrt{5})^n$$

$$(2-\sqrt{5})^n = \binom{n}{0} 2^n - \binom{n}{1} 2^{n-1} \sqrt{5} + \binom{n}{2} 2^{n-2} (\sqrt{5})^2 - \dots$$

$$+ = 2 \binom{n}{0} 2^n + 0 + 2 \binom{n}{2} 2^{n-2} (\sqrt{5})^2 + 0 + \dots$$

$$② P(x) = (x^2 - 49) + (3x - 7)(-x - 7) + 2(2x + 14)$$

$$\begin{aligned}P(x) &= (\cancel{x+7})(\cancel{x-7}) \quad \downarrow \\&= (\cancel{x+7}) [x-7 - (3x-7) + 7]\end{aligned}$$

$$-x-7 = -(\cancel{x+7})$$

$$⑤ \sum_{k=0}^n \left(\sum_{j=k}^n 2^j \right) = \sum_{k=0}^n (2^k + 2^{k+1} + \dots + 2^n)$$

$$1 + x + \dots + x^n = \frac{x^{n+1} - 1}{x - 1} \quad \text{qd xp1}$$

$$\cancel{\frac{2^{n+1} - 1}{2 - 1}}$$

$$2^k \left(1 + 2 + 2^2 + \dots + 2^{n-k} \right) = 2^k \left(\frac{2^{n-k+1} - 1}{2 - 1} \right)$$

Equa Diff d'ordre 1

1] Rappel de calcul intégral

1.1. Primitives usuelles

$$\int x^n dx = \frac{x^{n+1}}{n+1} + \text{cte} \text{ pour } n \neq -1$$

$$\text{Car } \begin{cases} (x^a)' = ax^{a-1} \\ (x^{n+1})' = (n+1)x^n \end{cases}$$

$$\int \frac{1}{x} dx = \ln|x| + \text{cte}$$

$$\Rightarrow \int \frac{u'}{u} = \ln|u| + \text{cte}$$

$$\int e^x dx = e^x + \text{cte}$$

$$\int e^{2x} dx = \frac{e^{2x}}{2} + \text{cte}$$

$$\int \cos x dx = \sin x + \text{cte}$$

$$\int \cos(2x) dx = \frac{\sin(2x)}{2} + \text{cte}$$

$$\int \sin x dx = -\cos x + \text{cte}$$

12. Integration per parte (I&II)

$$\boxed{\int u'v = [uv] - \int uv'}$$

ex1: $\int_2^{37} \arctan(t) dt = I ?$ $\arctan(t)' = \frac{1}{1+t^2}$

$$I = \int_2^{37} 1 \cdot \arctan(t) dt$$

$\uparrow u'$ $\uparrow v$

I&II $\begin{cases} u'=1 & u=t \\ v=\arctan t & v'=\frac{1}{1+t^2} \end{cases}$

$$\begin{aligned} I &= \left[t \cdot \arctan(t) \right]_2^{37} - \int_2^{37} t \cdot \frac{1}{1+t^2} dt \\ &= 37 \arctan(37) - 2 \arctan 2 - \int_2^{37} \frac{t}{1+t^2} dt \end{aligned}$$

$$I = 37 \arctan(37) - 2 \arctan(2) - \frac{1}{2} \int_2^{37} \frac{2t}{1+t^2} dt$$

$$= " - \frac{1}{2} \left[\ln(1+t^2) \right]_2^{37}$$

$$= " - \frac{1}{2} (\ln(1+37^2) - \ln(1+2^2))$$

Ex2: $I = \int x \sin(2x) dx$ I22 $\begin{cases} u = x & u' = 1 \\ v' = \sin(2x) & v = -\frac{\cos(2x)}{2} \end{cases}$

$$I = \left[-\frac{x \cos(2x)}{2} \right] - \int 1 \cdot -\frac{\cos(2x)}{2} dx = -\frac{x \cos(2x)}{2} + \frac{1}{2} \int \cos(2x) dx$$

$$I = -x \frac{\cos(2x)}{2} + \frac{1}{2} \frac{\sin(2x)}{2} + \text{cote} = \frac{\sin(2x) - 2x \cos(2x)}{4} + \text{cote}$$

$$Cx3 \quad C = \int \cos x e^x dx$$

$$I\&I \quad \begin{cases} u' = \cos x & u = \sin x \\ v = e^x & v' = e^x \end{cases}$$

$$C = [\sin x e^x] - \int \sin x e^x dx$$

$$I\&I \quad \begin{cases} u' = \sin x & u = -\cos x \\ v = e^x & v' = e^x \end{cases}$$

$$C = \sin x e^x - \left(E[\cos x] - \int -\cos x e^x dx \right)$$

$$C = \sin x e^x + \cos x e^x - \int \cos x e^x dx$$

$$C = \sin x e^x + \cos x e^x - C$$

$$\Rightarrow 2C = \sin x e^x + \cos x e^x \Rightarrow$$

$$C = \frac{\sin x e^x + \cos x e^x}{2}$$

$$C = \frac{(\sin x + \cos x) e^x}{2}$$

1.3 Changement de variable

ex1: $I = \int \frac{2t}{1+t^2} dt$ changement : $u = 1+t^2$

$$u' = \frac{du}{dt} = 2t \Rightarrow du = 2t dt$$

$$\Rightarrow dt = \frac{du}{2t}$$

$$I = \int \frac{2t}{u} \frac{du}{2t} = \int \frac{1}{u} du = \ln|u| + C = \ln(1+t^2) + C$$

ex2: $I = \int_0^{\pi/2} t \cos(t^2) dt$ dt $u = t^2$ $\frac{du}{dt} = 2t$

Δ les bornes changent : qd $t=0, u=0$ et qd $t=\frac{\pi}{2}, u=\frac{\pi^2}{4}$ $du = 2t dt \Rightarrow dt = \frac{du}{2t}$

$$I = \int_0^{\pi/2} t \cos(u) \frac{du}{2t} = \frac{1}{2} \int_0^{\pi^2/4} \cos(u) du = \frac{1}{2} [\sin(u)]_0^{\pi^2/4}$$

$$I = \frac{1}{2} (\sin(\frac{\pi^2}{4}) - \sin(0)) = \frac{1}{2}(1-0) = \frac{1}{2}$$

ex3: $k = \int_{t=2}^{t=3} \frac{1}{t \ln(t)} dt$

chgt $u = \ln(t)$

$\frac{du}{dt} = \frac{1}{t} \Rightarrow dt = t du$

$$k = \int_{u=\ln(2)}^{u=\ln(3)} \frac{1}{u} \cancel{\times} du = \left[\ln(u) \right]_{\ln(2)}^{\ln(3)} = \underline{\ln(\ln(3)) - \ln(\ln(2))}$$

↗
qd $t=2 \quad u=\ln(2)$

II Equations Différentielles Linéaires

②.1 L'équation homogène

On intéressé à : $a(x)y'(x) + b(x)y(x) = 0$

sols: $y_H = A e^{-\int \frac{b}{a} dx}$