



FIG. 1.

Bursting bubbles

Henri Lhuissier¹ and Emmanuel Villermaux^{1,2}

¹IRPHE, Aix-Marseille Université, 13384 Marseille Cedex 13, France

²Institut Universitaire de France, France

(Received 28 May 2009; published online 11 September 2009)

[doi:10.1063/1.3200933]

A young bubble is a piece of bare spherical liquid shell of radius R , with uniform thickness h , formed from an air volume rising underneath a water pool. It bursts by nucleating a hole, opening at the constant tangential velocity $V = \sqrt{2\sigma/\rho h}$ balancing inertia with surface tension forces, and collecting liquid in its rim. The centripetal acceleration $\gamma = V^2/R$ exerted on the rim perpendicular to the shell surface induces its Rayleigh–Taylor destabilization whose wavelength $\lambda_{\perp} \sim \sqrt{\sigma/\rho\gamma}$ scales as the geometrical mean of the only two length scales characterizing the bubble,

$$\lambda_{\perp} \sim \sqrt{Rh}. \quad (1)$$

Examples shown here include: [Fig. 1(a)] a snapshot of the rim instability and ligaments expulsion from the full

bursting sequence [Fig. 1(b)] of a $R=10$ mm radius bubble. $\Delta t=2$ ms between frames. Two water bubbles with distinct thicknesses h lead to different wavelengths λ_{\perp} : [Fig. 1(c)] $R=14$ mm, $\Delta t=5$ ms between frames and $\lambda_{\perp}=2$ mm; and [Fig. 1(d)] $R=11.5$ mm, $\Delta t=1.33$ ms between frames and $\lambda_{\perp}=0.8$ mm.

Ligaments emerge from the rim unstable crests and are stretched by centrifugation. They ultimately break by a capillary instability setting the resulting drop size distribution in the spray, as seen in Fig. 1(e). The distribution width is fixed by the relative ligament radius corrugations.¹ The drops thus formed are called “film drops.”² They originate in nature from bubbles entrained below breaking waves,³ and are believed to contribute significantly to the net water evaporation from the sea.⁴

¹E. Villermaux, “Fragmentation,” *Annu. Rev. Fluid Mech.* **39**, 419 (2007).

²D. C. Blanchard and L. D. Sysdek, “Film drop production as a function of bubble size,” *J. Geophys. Res.* **93**, 3649, DOI:10.1029/JC093iC04p03649 (1988).

³G. B. Deane and D. Stokes, “Scale dependence of bubble creation mechanisms in breaking waves,” *Nature (London)* **418**, 839 (2002).

⁴E. L. Andreas, J. B. Edson, E. C. Monahan, M. P. Rouault, and S. D. Smith, “The spray contribution to net evaporation from the sea: A review of recent progress,” *Boundary-Layer Meteorol.* **72**, 3 (1995).