

Introduction to Quantum Programming and Semantics


Louis LEMONNIER



THE UNIVERSITY *of* EDINBURGH
informatics

11th February 2025. Lecture 9: Complementarity

Previously



- Complementary bases
- Complementary Frobenius structures
- Equivalence
- Bialgebra

Complementary bases

Complementary bases: definition

$$\mathbb{C}^2 \quad \left\{ \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \end{pmatrix} \right\} \quad \left\{ \begin{pmatrix} 1/\sqrt{2} \\ 1/\sqrt{2} \end{pmatrix}, \begin{pmatrix} 1/\sqrt{2} \\ -1/\sqrt{2} \end{pmatrix} \right\}$$

Def two bases $\{a_i\}, \{b_j\}$ of a Hilbert space H
they are complementary if there exists $s \in \mathbb{R}$ constant
s.t. for all i, j $\langle a_i | b_j \rangle \langle b_j | a_i \rangle = s$.

Complementary bases: examples

\mathbb{C}^2

$$\left\{ \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \end{pmatrix} \right\}$$

$$\left\{ \begin{pmatrix} 1/\sqrt{2} \\ 1/\sqrt{2} \end{pmatrix}, \begin{pmatrix} 1/\sqrt{2} \\ -1/\sqrt{2} \end{pmatrix} \right\}$$

$$\left\{ \begin{pmatrix} 1/\sqrt{2} \\ i/\sqrt{2} \end{pmatrix}, \begin{pmatrix} 1/\sqrt{2} \\ -i/\sqrt{2} \end{pmatrix} \right\}$$

eigenvectors of respectively

$$Z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

$$X = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

$$Y = \begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix}$$

(Pauli gates.)

Complementary Frobenius structures

Complementary Frobenius structures: definition

 two Frobenius structures.

Def They are complementary if;



Special case:



Complementary Frobenius structures: example

$$\begin{array}{c} \text{hook} \end{array} \hat{=} \begin{array}{c} \text{cup} \end{array} \quad \text{cap} \hat{=} \text{C}$$

$$\begin{array}{c} \text{hook} \end{array} \hat{=} \begin{array}{c} \text{cup} \end{array} \quad \text{cap} \hat{=} \alpha$$



$$= \text{cup} \circ \text{cap}$$



Characterisation with unitaries

Prop Two Frobenius structures are complementary iff

 is unitary.

reminder: $(\text{---} \circlearrowleft)^{\dagger} = \text{---} \circlearrowright$ $(\text{---} \circlearrowright)^{\dagger} = \text{---} \circlearrowleft$ $(\text{---} \text{---})^{\dagger} = \text{---} \text{---}$

proof.



Equivalence

Th. In Hilbert spaces,

- compl. bases with $s = 1$
- compl. Fröb. struct.

are equivalent.

Proof

\triangleleft_a copy-able for $\langle _ \rangle$

\triangleleft_b copy-able for $\langle _ \rangle$

$$\langle b|a \rangle \langle a|b \rangle = \begin{array}{c} b \\ \triangleleft \triangleright a \\ a \triangleright \triangleleft b \end{array} =$$




$$\triangleleft_a = \triangleleft_a \langle _ \rangle$$



Further notions

Bialgebra and Hopf algebra


Bi algebra :



Hopf algebra :



Qubit gates

 $|0\rangle = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ $|1\rangle = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$

 $|+\rangle = \begin{pmatrix} 1/\sqrt{2} \\ 1/\sqrt{2} \end{pmatrix}$ $|-\rangle = \begin{pmatrix} 1/\sqrt{2} \\ -1/\sqrt{2} \end{pmatrix}$

CNOT $\hat{=}$ 

 = 



- Different bases in Hilbert spaces can be complementary
- Equivalent definition through bases or diagrams
- This complementary is necessary to represent quantum operations